



Review

Comprehensive review of the basic chemical behaviours, sources, processes, and endpoints of trace element contamination in paddy soil-rice systems in rice-growing countries

Waqar Ali^{a,b}, Kang Mao^a, Hua Zhang^{a,*}, Muhammad Junaid^c, Nan Xu^c, Atta Rasool^e, Xinbin Feng^a, Zhugen Yang^d

^a State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Guiyang 550081, China

^b University of Chinese Academy of Sciences, Beijing 100049, China


^c Key Laboratory for Heavy Metal Pollution Control and Reutilization, School of Environment and Energy, Peking University Shenzhen Graduate School, Shenzhen 518055, China

^d Cranfield Water Science Institute, Cranfield University, Cranfield MK43 0AL, United Kingdom

^e Department of Environmental Sciences, COMSATS University, Islamabad Vehari Campus, Vehari 61100, Pakistan



GRAPHICAL ABSTRACT



ARTICLE INFO

Editor: OK Yong Sik

Keywords:

Trace metal contaminants

Paddy soil system

Health effects

Transportation mechanisms

Accumulation

ABSTRACT

Rice is the leading staple food for more than half of the world's population, and approximately 160 million hectares of agricultural area worldwide are under rice cultivation. Therefore, it is essential to fulfil the global demand for rice while maintaining food safety. Rice acts as a sink for potentially toxic metals such as arsenic (As), selenium (Se), cadmium (Cd), lead (Pb), zinc (Zn), manganese (Mn), nickel (Ni), and chromium (Cr) in paddy soil-rice systems due to the natural and anthropogenic sources of these metals that have developed in the last few decades. This review summarizes the sources and basic chemical behaviours of these trace elements in the soil system and their contamination status, uptake, translocation, and accumulation mechanisms in paddy soil-rice systems in major rice-growing countries. Several human health threats are significantly associated with these toxic and potentially toxic metals not only due to their presence in the environment (*i.e.*, the soil, water, and air) but also due to the uptake and translocation of these metals *via* different transporters. Elevated concentrations of these metals are toxic to plants, animals, and even humans that consume them regularly, and the uniform deposition of metals causes a severe risk of bioaccumulation. Furthermore, the contamination of rice in the global rice trade makes this a critical problem of worldwide concern. Therefore, the global consumption of contaminated rice causes severe human health effects that require rapid action. Finally, this review also summarizes the available management/remediation measures and future research directions for addressing this critical issue.

* Corresponding author.

E-mail address: zhanghua@mail.gyig.ac.cn (H. Zhang).

1. Introduction

The word “paddy” is taken from the Malay word “padi,” which means “rice plant”; paddy soil, a well-known soil type, is submerged during rice cultivation (ar Hashmi and Varma, 2018). Trace metal contamination in paddy soil-rice systems poses a severe threat to the production of safe rice crops worldwide (Arunakumara et al., 2013). Trace element contamination in terrestrial resources is a focus of various environmental studies and requires more attention worldwide due to the toxicity, persistence, and bioaccumulation of these elements (Zhao et al., 2010; Lu et al., 2018). Agricultural soil has been contaminated by untreated waste discharge, emissions from rapidly intensifying industrial areas, increasing mining activities, chemical fertilizer use, industrial effluent disposal practices, and domestic wastewater (Lu et al., 2015). Trace metals such as As, Se, Cd, Pb, Zn, Mn, Ni, and Cr constitute an imprecisely defined group of inorganic biochemical hazards that are mainly found at polluted sites (Ali et al., 2019a; Garba and Abubakar, 2018). Soils are a substantial sink for trace metals released in unconfined environmental matrices due to human activities. Unlike organic pollutants, trace metals are not oxidizable to carbon dioxide (CO₂) through microbial action. Trace metals do not undergo biochemical degradation (Olowookere et al., 2018), and they persist in the soil system for a long time once they are released into the environment (Liu et al., 2013a). The presence of trace metals in paddy-soil systems can severely constrain the biodegradation of organic pollutants (Wuana and Okieimen, 2011).

The excessive concentrations of trace elements in agricultural soil systems result in increased trace element uptake by food crops such as paddy rice, wheat, maize, and vegetables (Rinklebe et al., 2019). Trace element contamination in paddy soil-rice systems poses severe threat to the human population as well as to ecosystems via direct ingestion or interaction with polluted soils and contaminated water; these contaminants become a part of food chains, such as the paddy soil-rice-human and paddy soil-rice plant-animal-human food chains. The rapid increase in the human population has led to a decrease in food quality and food safety due to the attenuation of agricultural lands, which has triggered food uncertainty and agricultural land occupancy problems (Wuana and Okieimen, 2011). Unpolluted agricultural areas are polluted with trace metals from polluted water that comes directly from cities or from other anthropogenic sources and is used for irrigation purposes; the constant use of agro-chemicals and fertilizers is another source of trace metals (Nieder et al., 2018; Geng et al., 2019). Such paddy soils expose humans to a high level of trace elements, and children are at a significantly higher health risk than adults according to risk assessment indices (Rinklebe et al., 2019).

Paddy cultivation in this respect is of extreme concern since crops need a very high quantity of irrigated water from seed growth to crop harvesting (Palaniappan and Annadurai, 2018). Approximately 160 million hectares of the global agricultural area is under rice cultivation, and rice is the leading staple food for more than half of the world's population (Seck et al., 2012). The category of total agrarian land includes paddy rice cultivation, and the average paddy rice production has progressively increased from 1.8 metric tons/hectare in 1960 to 4.3 metric tons/hectare currently (Arunakumara et al., 2013). Over the last few decades, rice consumption has increased, which has made it a valuable product (Timmer, 2014). Approximately 92 % of the total world paddy rice is produced in Asia, and paddy rice is a well-known staple food consumed by most of the world's population (Wang et al., 2018). Trace metal pollution in the paddy-soil system and its successive accumulation in different parts of the rice plant (roots, shoots, leaves, and grains) have been observed in different rice-growing countries (Hu et al., 2016). Hence, trace metal contamination in paddy soils and its transfer in paddy soil-rice systems are a growing concern (Halim et al., 2015). Several studies in various countries have investigated certain specific trace elements, such as As, Se, Cd, Pb, Zn, Mn, Ni, and Cr (Rinklebe et al., 2019).

Currently, measuring or predicting the likely human health risks of trace elements that primarily originate from exposure to metal-polluted soil, water, and atmospheric deposition is becoming more common (Antoniadis et al., 2019a). These trace elements are mainly studied in works evaluating the potentially toxic trace elements through human health risk assessments specifically targeting children and male and female adults (Abraham et al., 2018). The human health risks of metal-polluted soils can be measured by using the soil pollution indices of trace elements, soil to plant transfer factors, and their concentrations in rice systems (Antoniadis et al., 2019a; Palansooriya et al., 2020). Recent studies have stated that trace element accumulation in paddy soil systems is related to risks to soil fertility and quality, as well as soil biochemical activity such as microbial and enzyme activity. Elevated trace element concentrations in paddy soil systems are primarily known to affect the microbial count in soil and the associated microbial activities, which directly influence the fertility of the soil (Minnikova et al., 2017). Furthermore, the public is becoming more conscious of the implications of trace element contamination in agricultural soils and its severe toxic effects on human and environmental health as well as of the enhancement and expansion of technologies for trace element removal/clean-up at polluted sites (Liu et al., 2018). Hence, it is imperative to deploy recent advances and various site-specific contamination clean-up technologies that might feasibly and effectively remediate trace element pollution in soils. In addition, a summary of recent review papers about trace element contamination in soil-plant and rice system topics, objectives and gaps is discussed in Table S1.

The present study aims to systematically review and understand the basic behaviours of trace metals in the soil system to assess sources, uptake, translocation and accumulation mechanisms, the status of trace element contamination, and associated risks in paddy soil-rice systems in rice-growing countries and contaminated soil management/remediation methods.

2. Trace element contamination in soil: Possible sources

2.1. Natural sources

Trace metals persist naturally in the soil environment mainly due to pedogenic processes of parent material weathering at a concentration that is regarded as trace (< 1000 mg/kg) and are infrequently poisonous (Wuana and Okieimen, 2011). It is deceptive that the significant preliminary source of trace metals in the soil system is the parent material (Zhao et al., 2014a). The effect of parent materials in defining the trace element content is notable in soil systems, which have formed from the same or an individual parent material, such as igneous rocks, sedimentary rocks, and/or new soils weathered under temperate environments (Bini, 2013). As soon as the rock weathering process takes place, the main crystal-like structures of the raw materials are entirely broken down and allow trace metals to either be adsorbed on the soil surface or transported to the surface or groundwater system (Arunakumara et al., 2013). The effect of rock parent materials on the type and content of trace elements in the soil system are mainly reliant on pedogenic processes (White, 2013). However, trace metal concentrations in soil systems are primarily derived from pedogenic parent materials, and the physicochemical and biological features of soil determine the extent of trace element accumulation (Huot et al., 2015). Hence, trace element contamination in the soil system primarily results from the natural availability of extensive parent material (glacial till), and anthropogenic activities are also significant sources (Dung et al., 2013).

Accordingly, uncovering a strong association between trace element status in the soil system and the parent material is demanding (Chen et al., 2008). According to (Jun-Hui and Hang, 2009; Hu et al., 2019), a compilation of results from the monitoring of significant sources of trace elements in paddy soil and a pollution assessment of the Yangtze River Delta and Fuyang County in China, respectively, suggested that

the Ni concentration of the paddy-soil system was determined to be high due to the parent material and that Ni showed the most extreme high concentration of all contaminants.

Another study from Dongguan, China, described the status of trace element levels in agricultural soils and in different vegetables with diverse key sources. Cu, Zn, Ni, Cr, and As, were mainly derived from natural rock parent materials. Nevertheless, Cd and Pb were present due to anthropogenic activities and were above the recommended concentrations of trace metals in soil systems; this was not true of the parent material (Cai et al., 2010).

The critical relationship between mafic rock types and levels of trace elements such as Ni, Al, Mn, Mg, and V in the soil system suggests the significance of natural rock parent material to the spatial dispersal of these trace elements (Alloway, 2013a). Mafic and ultramafic rocks accumulate trace metals, including Cd, Cr, and Ni, leading to higher concentrations in soil-plant systems (Islam et al., 2016). In Pakistan, studies conducted by (Khan et al., 2010; Tariq, 2001) on soil and vegetable accumulation of trace metals from natural geological sources in Gilgit, northern Pakistan, and trace metals in the piedmont soils of the Peshawar Basin, Pakistan, respectively, suggested higher levels of trace elements, including Cd, Pb, Zn, Cu, Cr and Ni, in agricultural soils that were primarily released from parent rocks and calcite rocks. The release of Mn and As mainly from natural aquifer rocks has been reported in soil and groundwater systems in different locations in Bangladesh and West Bengal, India (Fendorf et al., 2010; Rahman et al., 2007). Nevertheless, although trace metals usually originate from natural sources, the type and elemental composition of parent material, the extent of the rock weathering process, the physicochemical and biological properties of soil, and the climatic environment seem to have a significant effect on the occurrence and distribution of trace elements in soil systems (Essington, 2015).

2.2. Anthropogenic sources

2.2.1. Agricultural activities

2.2.1.1. Intensive pesticide use. The term “pesticides” includes any combination of substances intended for preventing, destroying, resisting and mitigating any pests such as weeds, nematodes, insects, mites, and rats. Consequently, it comprises insecticides, nematicides, herbicides, fungicides, molluscicides, and other materials extensively used for pest control (Arunakumara et al., 2013). Despite the significant influence of pesticides on worldwide agricultural production, the adverse human health effects linked with them is causing increasing concern (Guedes et al., 2016). However, the intensive use of insecticides has grown extensively over the past decades, and in recent decades, approximately 2.3 million tons of pesticides have been used yearly (Zhang et al., 2011).

Extremely hazardous substances, such as persistent organic pollutants (POPs), organophosphates and organochlorine insecticides such as dichlorodiphenyltrichloroethane (DDT), chlordane, lindane, and chlorinated hydrocarbon endosulfan) are used in the agricultural application of different conventional pesticides reported to contain significant levels of trace elements and enter environmental matrices (Yadav et al., 2015); these pesticides are based on chemical mixtures comprising Cu, Ni, Cr, Mn, Cd, As, Pb, and Zn (Wuana and Okieimen, 2011). The Bordeaux mixture of copper sulfate (CuSO_4) is commonly used to control mycological attacks in several crop systems (Hajek and Eilenberg, 2018). Arsenic and As-containing compounds, such as lead arsenate (PbHAsO_4) insecticides, have been intensively used in the past to control pests in different crop systems (Gupta et al., 2017). However, the intensive use of fertilizers is also common practice to reach maximum agricultural production from a unit area of land (Savci, 2012). Agrochemical fertilizers usually contain significant concentrations of different trace metals and their metalloids, including As, Cd, Cr, Zn Cu, Mn, Ni, Zn, & Pb (Arunakumara et al., 2013). Radionuclide trace elements such as (U, Ra, & Th) have also been introduced into agricultural

soil along with fertilizers (Sahu et al., 2014). Nevertheless, some of the trace metals (Cu, Mn, Mo, Zn, Ni, & Fe) at low concentrations are considered essential nutrients (Alloway, 2013b). Agrochemical fertilizers contain significant levels of trace elements such as Ni, Pb, Fe, Co, & Cd as impurities (Arunakumara et al., 2013).

2.2.1.2. Intensive use of fertilizers. Worldwide, the extensive use of fertilizers for agricultural purposes has increased exorbitantly in the previous six decades. In 2006, the World Resources Institute (WRI) specified that five key countries, China, the United States, India, Indonesia, and Brazil, were responsible for two-thirds of all fertilizer consumption; China ranked first with approximately 32.9 % (55,925.6 thousand metric tons) consumption of total fertilizer worldwide (Gupta et al., 2014). Fertilizer usage in India has increased intensively to 133 kg/ha in 2011 from 1 kg/ha in 1951–1952 (Gupta et al., 2014). Among the various types of fertilizers, the application of phosphate-containing fertilizers is the most common. It is a dynamic factor in the sustained agricultural growth and high productivity of rural farming in several countries around the world (Sattari et al., 2014).

In Asia, the typical application of phosphate-based fertilizer is approximately 34 kg/ha (Gupta et al., 2014). However, the concentrations of pollutants in fertilizer vary significantly, especially the concentrations of trace metals. According to the literature, the application of phosphate-based fertilizer at a rate of 20 kg/h/year in soil leads to the introduction of approximately 25 g/h/year and 0.01 g/ha/year of Cr and Hg, respectively (Gupta et al., 2014). Phosphate-based fertilizers have been a significant source of trace metals compared with other inorganic fertilizers, as they may contain 0.1–170, 1–12, 7–38, & 7–225 mg/kg of the total Cd, Co, Ni, & Pb, respectively (Arunakumara et al., 2013; Ihedioha et al., 2016). Phosphate-based fertilizers can relocate naturally available As into agricultural soil systems, which causes it to accumulate in runoff (Ravenscroft et al., 2009). Agrarian soil in European countries can accumulate As at a rate of 7.7 g/ha/year, compared with Asian countries such as China, which can accumulate As at a rate of 2.5 g/ha/year through the extensive application of phosphate-based fertilizers (Arunakumara et al., 2013; Luo et al., 2009; Nziguheba and Smolders, 2008). Nevertheless, phosphate fertilization may reduce metal toxicity and mobility, as well as the plants available in polluted paddy soils and other agrarian soils (Shahid et al., 2014). A field trial study was conducted in metal-contaminated soil with mining tailings in Shaoxing, China, and suggested that the addition of phosphate-based fertilizers could obviously reduce the levels of water-extractable elements such as Zn (11.7–75.3 %), Pb (22.0–81.4 %), and Cd (1.5–30.7 %) (Wang et al., 2008).

In addition to phosphate-based fertilizers, mixed fertilizers also act as significant sources of trace metal contamination in agrarian soils (Wortman and Lovell, 2013). In China, the yearly received rates of Cd, Cr, Pb, and Ni from phosphate-based mixed fertilizers are 0.73, 28.1, 6.9, and 4.1 g/ha, respectively (Luo et al., 2009). In China, it has been reported that approximately 1200 tons of Zn and 5000 tons of Cu-containing fertilizers were added to agrarian lands per annum (Luo et al., 2009). Agrarian soil contamination by trace metals through fertilizer application has been found in several countries, such as China, India, Malaysia, Bangladesh, Brazil, and certain European countries (Arunakumara et al., 2013; Satpathy et al., 2014; Hezbullah et al., 2016).

Nutritionally rich wastewater provides crops with essential as well as toxic trace metals and accumulates in agrarian soil with organic matter (Wang et al., 2017). Despite its inherent quality and shortcomings, the application of wastewater for irrigation purposes has been widely practised for many decades worldwide (Jiménez et al., 2009). Primarily, in arid zones around the world, where water is a limiting factor for agrarian growth, urban wastewater is often used as a consistent source of irrigation for crop cultivation (Clarke, 2013; Pettygrove, 2018). The wastewater is usually applied directly for irrigation and circumvents the essential requirements for wastewater

treatment plants; the accumulation of severely toxic trace elements in agrarian soils through consistent use of wastewater has often been observed in different countries (Elgallal et al., 2016). It has been estimated that approximately 20 million ha of land is irrigated with wastewater in several countries, including China, India, Bangladesh, Pakistan, Germany, and France (Arunakumara et al., 2013; Mateo-Sagasta et al., 2015; Biswas et al., 2017; Murtaza et al., 2010). Depending on the primary source, the wastewater might include different trace metals, such as As, Zn, Cd, Cu, Pb, Cr, Mn, or Ni, in varying concentrations (Baysal et al., 2013). However, it has also been reported that after going through wastewater treatment plants, trace metals can persist in wastewater and can accumulate in living systems through food chains (Rajasulochana and Preethy, 2016).

Hence, before the reuse of wastewater for agrarian purposes, significant consideration is necessary for public health as well as acceptance (Qadir et al., 2010). In an eastern suburb of Beijing, China, a study reported elevated concentrations of trace metals in wastewater-irrigated soils to be in the order of Cr (78.3) > Zn (71.5) > Ni (31.8) > Cu (25.5) > Cd (0.81) mg/kg; however, in wastewater-irrigated agrarian topsoils compared with Chinese reference soils, Cr (53.8) > Zn (51.9) > Ni (18.7) > Cu (16.0) > Cd (0.36) mg/kg, which are within the World Health Organization (WHO) and the national standards of China, except for Cd (Arunakumara et al., 2013; Chen et al., 2009a). A similar type of study was carried out in the Zhangshi wastewater irrigation area in a suburb of Shenyang City, China, in an agrarian area with about three decades of irrigation history. The elevated trace metal-containing wastewater and its measured concentrations were found to be in the order Ni (241.6) > Pb (88.1) > Cu (69.2) > Cd (2.7) mg/kg; the levels were significantly elevated in wastewater-irrigated topsoil compared with those in reference soils (Zhang et al., 2008).

A study conducted at the Bindal River, Dehradun, India found that wastewater irrigation caused a higher percent concentration of trace metals in agrarian soil, in the order Zn (48 %) > Pb (20 %) > Ni (13 %) = Cu (13 %) > Cr (5 %) > Cd (1 %), which exceeded the permissible limits defined by the WHO and by India (Pathak et al., 2010). Though the agrarian soil was irrigated with treated wastewater in Varanasi, India, the concentrations of trace metals were found in the following

order: Pb (123.5) > Zn (122.3) > Cu > (77.9) > Cr (56.3) > Cd (3.9) mg/kg; these levels exceed the limits of both the WHO and India (Mishra and Tripathi, 2008). A study conducted in Abbottabad, Pakistan, reported on the status of trace metal concentrations in municipal wastewater used for agrarian purposes, including Cd (2.8 mg/kg), Cr (4.2), Cu (28), Ni (7.5), Zn (135.6), Pb (15.6) and Fe (270) mg/kg, which are below the standards defined by the USA, UK and EU (Hassan et al., 2013). However, a study conducted in Hanoi, Vietnam, in a paddy soil system found that the irrigation wastewater was contaminated with wastes from several industrial plants and reported the average concentrations of trace metals, such as Cu (202), Zn (192), Pb (159), and Cd (4) mg/kg, which all exceeded the WHO and Vietnamese defined standards (Huong et al., 2008). Sewage sludge or biosolids are mainly reliable organic products produced by wastewater treatment and can be constructively recycled (Cimen et al., 2014).

Different waste disposal approaches, such as agricultural application, landfilling, incineration, and ocean dumping, are used based on resource accessibility (Yin et al., 2018). Agricultural practices that use sewage sludge are widely known as an economically feasible and ecologically acceptable method of disposal (Chen et al., 2012). Therefore, an extensive quantity of sewage sludge is directly deposited on agrarian lands in China, where approximately 10 % of the total sewage sludge produced in the country is applied directly to agricultural land (Yang et al., 2015). From an agrarian perspective, the direct use of sewage sludge is considered to be an excellent fertilization method because it can improve the physiochemical and biological properties of the soil and contains high levels of nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), and other trace elements that are essential for crop growth (Singh et al., 2011a; Głab et al., 2018). Nevertheless, elevated concentrations of trace metals such as Cu, Zn, Cr, As, Se, Cd, Mn, and Pb, are often reported in sewage sludge and may cause adverse effects on humans and the environment (Babayemi et al., 2016; Singh and Agrawal, 2010). Trace metal contamination in agrarian soils from the application of sewage sludge has been reported as one of the primary concerns about this method because of the long-term accumulation of trace metals in the soil-plant system (Guoqing et al., 2019). Obviously, the trace metal content in sewage sludge differs by numerous factors, such as the quality of wastewater and the effectiveness and




Fig. 1. A general overview of natural, anthropogenic sources of trace elements in environmental matrices, and associated human health toxicities.

meticulousness of the treatment process (Nair and Ahammed, 2015). Usually, the contents of trace elements in sewage sludge can vary from 0.5 to 2 % on a dry basis (Arunakumara et al., 2013). However, the frequent application of livestock manure in agrarian soil is recognized as a vital source of trace metals in soil crop systems and causes various environmental issues and human health effects (Arroyo et al., 2014; Hu et al., 2017).

Livestock manure is a valuable source of organic compost that can improve paddy soil fertility as well as the organic matter content (Zhang et al., 2018). Livestock are also frequently fed with several feed additives to improve growth performance and reduce the risk of human disease (Hu et al., 2017; Zain, 2011). However, feed additives and animal manure are significant sources of trace metals at various concentrations (Hu et al., 2017; Ziarati et al., 2018). The continuous application of livestock manure to agrarian soil causes the accumulation of trace metals in the soil crop system and food chain, which seriously threatens the environment, the quality of plant crop production, and human health (Zhao et al., 2014b).

2.2.2. Industrial activities

Mining activities provide several socio-economic benefits but are considered some of the most hazardous activities worldwide (Arunakumara et al., 2013; Zhengfu et al., 2010). Mining activities such as grinding, concentrating minerals, and tailings and mine and mill wastewater disposal are significant and confirmed sources of trace metal contamination in soil crops (Fig. 1), mainly in paddy soil-rice systems (Krishna et al., 2013; Wang et al., 2016). Trace metal contamination in paddy soil-rice systems through mining activities is considered a significant issue worldwide (Halim et al., 2015). The spatial distributions of trace metal concentrations adjacent to mining sites depend on geochemical conditions, and mineralization following the release of untreated wastes into the environment may increase trace metal concentrations in agrarian soil and food crop systems (González-Corrochano et al., 2014; Candeias et al., 2014). According to the literature, a study conducted near an abandoned copper mine in eastern China suggests that continuous mining practices may increase trace metal contamination in soils, with average metal concentrations of Cu (147), Pb (53.8), Zn (158), and Cd (0.32) mg/kg (Qin et al., 2012).

A study conducted in paddy soils adjacent to a mining site in the Kočani field, Macedonia, was severely affected by an increased level of trace metals such as As (47.6), Cd (6.4), Cu (99), Pb (983) and Zn (1245) mg/kg because of the wide dispersal of trace metals from metal mining activities (Rogan et al., 2009). However, a similar type of study conducted in Vietnam in paddy fields adjacent to chromite mine tailings piles showed trace metal concentrations of Co (375), Ni (5590), and (5750) mg/kg (Kien et al., 2010). Another study was conducted in Thai Nguyen Province in northern Vietnam in paddy fields 50 m away from a Pb recycling smelter and reported significantly elevated concentrations of trace metals, including Pb and Cd, ranging from 1100 to 7000 and 0.5 to 1.3 mg/kg, respectively (Chu, 2011).

In Hezhang County in China, smelting practices commonly cause Cr pollution ranged from 71 to 240 mg/kg in agrarian soils (Bai et al., 2009). A study around the chromite mining site at Chikkondanahalli in the Nuggihalli schist belt, Karnataka, India, found that the status of trace metals in agrarian soil concentrations was in the order Cr (791) > Ni (168.5) > Co (21.5); these levels exceed permissible soil quality limits and suggest that the elevated levels of these elements are correlated with previous and current uncontrolled mining activities in the study area (Krishna et al., 2013). Airborne fine-grained fly ash waste, another primary source, contributes to trace metal contamination in environmental matrices (Funari et al., 2016). The reduction in petroleum-based sources causes increasing consideration of renewable energy production, primarily through biomass, including wood, straw, and crop burning (Mushtaq et al., 2014). However, coal-based thermal power generation plants are still a well-known energy source in various countries, such as India, China, Pakistan, Japan, and Korea (Awan and

Rashid, 2012; Guttikunda and Jawahar, 2014). Thus, the production of incinerated airborne fine-grained solid fly ash waste is considered a serious concern worldwide (Tao et al., 2015). Substantial amounts of fly ash containing high concentrations of trace metals are produced through the incineration process (Tao et al., 2015).

Moreover, fly ash neutralizes agrarian soil acidity, improves soil structure and texture, reduces bulk density, and increases soil water and nutrient-holding capabilities, suggesting the rationality of its use in crop production (Arunakumara et al., 2013). Therefore, the direct use of fly ash in agricultural lands is a conventional practice that increases trace metal concentrations and bioaccumulation in crops, mainly in paddy soil-rice systems (Nayak et al., 2015). The direct disposal of hazardous waste has also been reported as a significant contributor to elevated levels of trace metals in soil-crop systems (Wuana and Okieimen, 2011). The distributions of trace metals from hazardous waste disposal sites to adjacent areas depend on various interconnected geochemical activities (Bretzel and Calderisi, 2011). The distributions of trace metals to adjacent agricultural soils are usually influenced by soil pH, oxidation-reduction potential (ORP), and the availability of involved agents such as carbonates, sulfates, chlorides, microbial activities, organic, and hazardous waste properties (Parth et al., 2011; Bolan et al., 2014).

A study was conducted in Dholpur district, Rajasthan, India, and reported trace metals, including Cr, Cu, Fe, Ni, Pb, and Zn, with elevated concentrations at dumpsites (34.56 ± 1.05 – 110.75 ± 2.59 mg/kg) (Jadoun and Singh, 2014). A similar study was conducted adjacent to dumping sites in Allahabad, Uttar Pradesh, India, and proposed that the soil pH was negatively correlated with Ni and Fe, though the contents of Zn were significantly positively correlated with Pb and Cu (Tripathi and Misra, 2012). However, in the northwest region of Hyderabad, India, several hazardous waste dumping sites exist where approximately 50,000 tons of hazardous/industrial waste is abandoned yearly in landfills, which causes elevated trace metal contamination in adjacent soil, including Cr (127.9), As (51.7) and Pb (206.4) mg/kg (Parth et al., 2011).

The increase in the global population has increased the use of electronic appliances, which cause 20–50 million tons of annual production of e-waste worldwide, which is a serious concern (Borthakur and Govind, 2017; Zhang et al., 2012b). E-waste can also cause pollution from numerous trace metals. (Tue et al., 2016). The reviewed status of trace metal contamination in soil systems due to e-waste in China comprises several study results suggesting Cu and Cd concentrations of thousands of mg/kg beyond the permissible limits of 50 and 0.3 mg/kg for agrarian soils (Zhang et al., 2012b).

However, in a study conducted in Taizhou, China, paddy fields adjacent to an e-waste recycling zone contained trace metal concentrations of 61.21, 98.81, 1.19, and 55.81 mg/kg for Cr, Cu, Cd, and Pb, respectively, which were beyond the background values (Fu et al., 2008). Another similar study conducted in southeastern Zhejiang Province in China investigated paddy fields near an e-waste recycling areas and found trace metal concentrations of 20.95, 97.90, 298.31, 0.62, and 44.29 mg/kg for Cr, Cu, Zn, Cd, and Pb, respectively, which were far beyond the background values (Jun-Hui and Hang, 2009).

3. Trace element basic chemistry, behaviour in the soil system, and associated risks

Although trace metals play a significant role in crop production, they may reduce crop production because of the risk of bioaccumulation and biomagnification in the food chain (Majed et al., 2016). A basic knowledge of trace metal chemistry, transport, environmental fate, and related human health effects is required to understand their speciation and bioavailability as well as the remediation options (Wuana and Okieimen, 2011).

The transport and fate of trace metals in soil crop systems mainly depend on the bioavailability, chemical form, and speciation of trace

Table 1
Reported trace element concentrations ranges minimum-maximum (mean) mg/kg in paddy soils of different countries.

Country	Locations	Cd	Cu	Ni	Pb	Zn	Cr	Mn	Se	As	Reference
China	Wenling	0.11–3.45 (0.31)	15.8–160.1 (41.1)	9.2–68.2 (33.9)	27.1–140.5 (48.30)	64.97–276.0 (137.0)	-	-	-	-	Zhao et al. (2010)
	Shengzhou	0.04–0.28 (0.1)	15.1–78.1 (29.8)	5.1–62.7 (19.2)	22.0–50.4 (34.3)	34.7–113.6 (75.8)	15.1–78.1 (29.8)	-	-	-	Zhao et al. (2009a,b)
China	Nanxun	0.12–0.78 (0.2)	18.8–82.3 (31.1)	18.5–55.6 (32.1)	20.2–51.7 (33.2)	61.5–266.9 (106.8)	-	-	-	-	Zhao et al. (2015b)
	Kunming	0.20–3.57 (1.1)	45.4–2026 (152.7)	24.1–95.8 (45.0)	6.8–146.6 (42.2)	55.8–484.9 (203.3)	(52.0–105.2 (73.4)	-	(10.0–141.5 (24.4)	(7.3–124.0 (19.5)	Cheng et al. (2018) Lu et al. (2018)
China	Hunan	0.17–27.3 (1.1)	-	-	24.8–266.0 (52.9)	-	-	-	-	-	Zeng et al. (2011)
	Nanhu	-	6.6–8.0 (7.3)	-	11.0–12.0 (11.4)	4.7–11.4 (7.7)	0.6–1.1 (0.8)	-	-	-	Zeng et al. (2015)
China	Tongxiang	-	5.26–9.70 (7.0)	-	9.1–17.8 (12.6)	2.26–6.94 (3.9)	0.5–0.7 (0.6)	-	-	-	Zeng et al. (2011)
	Xiaoshan	-	1.38–5.66 (3.7)	-	2.0–9.74 (5.1)	1.40–12.2 (5.7)	0.3–0.8 (0.6)	-	-	-	Zeng et al. (2015)
China	Shimen	0.06–0.4 (0.2)	-	26.6–48.7 (36.2)	19.8–32.3 (24.9)	-	23.2–33.5 (26.0)	-	-	3.16–45.2 (22.1)	Zeng et al. (2015)
	Fenghuang	0.04–28.0 (3.3)	-	25.9–44.5 (36.5)	19.6–83.6 (43.3)	-	21.5–28.2 (23.7)	-	12.2–34.8 (21.2)	3.3–6.3 (4.5)	Zeng et al. (2011)
China	Xiangtan	0.18–1.17 (0.5)	-	14.9–40.0 (27.7)	37.7–262.0 (94.5)	-	28.1–39.3 (33.1)	-	-	-	Zeng et al. (2011)
	Tan Long	2.3–42.9	-	-	1271–3953	-	-	-	-	-	Dang et al. (2008)
Bangladesh	Chi Dao	2.5–56.5	-	-	2000–10000	-	-	-	-	-	Chiu (2011)
	Faridpur	0.12–0.17 (0.2)	38.2–63.5 (48.4)	42.2–61.0 (48.8)	24.32–34.0 (26.8)	82.7–117.4 (97.2)	73.5–108.1 (86.0)	374.1–575.2 (449.7)	(0.6–1.4 (1.0)	17.4–65.0 (33.1)	Ahsan et al. (2009)
Bangladesh	Dhamrai	0.11–0.22 (0.2)	29.6–33.7 (31.8)	43.8–47.9 (46.1)	23.2–26.6 (24.8)	78.9–125.9 (98.8)	45.6–84.4 (76.1)	471.1–655.2 (553.7)	0.6–1.3 (1.0)	3.1–9.0 (6.1)	Khanam et al. (2019)
	several locations	0.40	-	-	4.90	-	-	-	0.32	3.6–26	Khanam et al. (2019)
India	Narora	0.6–4.6 (2.4)	6.41–153.5 (31.8)	-	2.0–46.0 (21.8)	-	3.9–21.5 (8.1)	3.3–35.5 (21.7)	-	-	Singh et al. (2014)
	several locations	(0.8)	-	-	0.90	-	-	-	0.23–4.52	61–416	Khanam et al. (2019)
India	Fatehabad	-	10.8–37.0 (21.8)	0.04–23.5 (10.4)	7.5–148.2 (54.4)	71.5–146.7 (106.0)	0.1–9.7 (6.5)	-	-	-	Yadav et al. (2017)
	Kota	(0.8)	-	-	-	-	21.10	46.40	-	-	Yap et al. (2009)
Japan	several locations	0.2–0.3	-	-	-	-	-	-	0.43	11.00	Khanam et al. (2019)
	several locations	0.45 ^a	-	39.0 ^a	29.0 ^a	99.0 ^a	64.0 ^a	-	-	-	Makino et al. (2010)
USA	Aomori	0.16–0.53 (0.37)	18.0–45.0 (29.0)	5.9–18.0 (11.0)	9.2–44.0 (21.0)	63.0–160.0 (110.0)	16.0–80.0 (31.0)	370.0–1500 (700.0)	0.72–1.4 (1.0)	6.7–34.0 (12.0)	Tsukada et al. (2007)
	several locations	(0.3)	-	-	(55.0)	-	-	-	-	15–950	Khanam et al. (2019)
Korea	several locations	(0.1)	-	-	(5.3)	-	-	-	0.1–1.1	0.78	Sun et al. (2010) and Dhillon and Dhillon (2018)
	several locations	0.1–1.01 (0.11)	0.1–41.6 (4.7)	-	0.01–66.4 (4.8)	0.1–96.7 (4.5)	-	-	-	-	Jo and Koh (2004)
Pakistan	Four mining areas	0.001–12.8 (2.3)	14.7–243.0 (63.5)	-	26.3–885.0 (146.0)	89.7–1600.0 (393.0)	-	-	-	2.8–704.0 (64.4)	Kwon et al. (2017)
	Shakargath	1.0–78.9 (35.0)	-	60.0–99.7 (85.8)	113.8–2967 (1065)	-	13.8–89.8 (37.8)	-	-	-	Tariq and Rashid (2012)
Thailand	Bangkok	0.01–0.07	0.7–3.0	0.4–2.6	0.2–1.4	1.0–14.4	0.1–1.9	-	-	-	Chinoim and Sinbathong (2010)

^a Denotes the total concentration.

metals (Islam et al., 2015). Once in the soil system, trace metals are absorbed through initial reactions that occur in minutes to hours, followed by slow adsorption reactions that occur in days to years; thus, their conversion and redistribution into numerous chemical forms with varying mobility, toxicity, and bioavailability take place (Huang et al., 2015). The dispersal and reactions of trace metals in the soil system are likely controlled by several factors, such as mineral dissolution and precipitation, metal ion exchange capacity, adsorption or desorption, biological mobilization or immobilization, soil-plant availability and uptake size and aqueous complexation (Table 1) (Wuana and Okieimen, 2011).

3.1. Arsenic

Arsenic mainly occurs in different compounds such as As_2O_3 , and can be recovered by the processing of ores usually containing Pb, Zn, Ag, Cu and Au (Rudra and Basher, 2020). There are four natural processes, i.e., reductive dissolution, sulfide oxidation, alkali desorption, and geothermal activities, involved in the release of As to the environment (Bhattacharya et al., 2007). Two main species of As exist in the terrestrial environment: inorganic pentavalent arsenic, arsenate As(V) under oxidizing (aerobic) conditions, and trivalent arsenite As(III) under reducing (anaerobic) conditions (Palansooriya et al., 2020; Sun et al., 2014). As(V) is dominant, primarily in the form of arsenate ions (AsO_4^{3-}) in different states, such as arsenic acid (H_3AsO_4), dihydrogen arsenate (H_2AsO_4^-), hydrogen arsenate (HASO_4^-) and arsenate ions (AsO_4^{3-}) (Wuana and Okieimen, 2011; Zhang, 2011). Thus, As and As(V) act as chelates and can easily precipitate once cations are present (Chiban et al., 2012). The As(V) complexes are stable under specific conditions and coprecipitate and absorb Fe-oxyhydroxide under reducing and/or partially reducing conditions. The coprecipitates are immobile under these particular environmental conditions; nevertheless, the mobility of As increases with an increase in pH (Campbell and Nordstrom, 2014; Wilson et al., 2010). However, under anaerobic conditions, As(III) is the dominant species of As, existing as an arsenite ion (AsO_3^{3-}) and its protonated forms, such as arsenious acid (H_3AsO_3), dihydrogen arsenite (H_2AsO_3^-) and hydrogen arsenite (HASO_3^{2-}) (Table S2) (Wuana and Okieimen, 2011). As(III) can coprecipitate or can adsorb with sulfides (S^{2-}) and has an affinity to bind with other sulfur compounds (Roy et al., 2015). As(III) and arsine (AsH_3) usually exist under highly anaerobic conditions (Wang et al., 2014). The main routes of human As exposure include drinking water and food (Wu et al., 2011). The bioaccumulation of As in living systems (plants & humans) takes place through methylation, which generates highly volatile methylated derivatives of AsH_3 compounds such as dimethyl arsine ($\text{HAs}(\text{CH}_3)_2$) and trimethyl arsine ($\text{HAs}(\text{CH}_3)_3$) (Rahman et al., 2012). Arsenic usually exists in anionic forms that do not exhibit an affinity to bind with anions, including chloride (Cl^-) and sulfates (SO_4^{2-}), to form complexes (Singh et al., 2015).

Arsenic can also exist in organometallic forms, which include methyl arsenic acid ($(\text{CH}_3)\text{AsO}_2\text{H}_2$) and dimethyl arsenic acid ($(\text{CH}_3)_2\text{AsO}_2\text{H}$) (Wuana and Okieimen, 2011). The modes of toxicity for As(III) and As(V) are different (Gailer, 2007). As(III) is sixty times more toxic and carcinogenic to humans than As(V) (Hughes et al., 2011). As(III) can bind with tissues for a longer period than As(V), which binds with specific groups of proteins that cause As(V) to act biologically against phosphate, distressing the production of ATP synthesis (Brown and Ross, 2002; Chandrakar et al., 2018). Long-term exposure to As damages the human cardiovascular, dermal, neurological, hepatic, respiratory, and reproductive systems (Fig. 1) (Ali et al., 2019b).

3.2. Selenium

Selenium exists in different oxidation states that permit the production of organoselenium and selenoamino acid complexes (Tinggi, 2003). In the soil system, selenate (SeO_4^{2-}) and selenite (SeO_3^{2-}) are

stable for extended periods (Hooda, 2010). Several Se(VI) compounds exist in soil, but only SeO_4^{2-} usually occurs in normal pH ranges, and the most dominant form of Se(IV) below soil pH 8.5 is hydrogen selenite (HSeO_3^-) (Hooda, 2010). Selenium is an essential trace element for animals and humans at a certain level; for adults, the recommended dietary allowance (RDA) of Se is 55 $\mu\text{g}/\text{day}$ (Sun et al., 2014; Zwolak and Zaporowska, 2012; Zeng et al., 2005), and an excessive amount of Se can cause severe toxicity. Se(IV) is commonly used as a feed additive in different commercial animal diets with a recognized Se dose of 0.5 mg/kg in the whole feed (Zwolak, 2019). In humans, Se intake varies across various countries (Ali et al., 2020).

Overall, Se consumption for adults ranges from 93 to 134 $\mu\text{g}/\text{day}$ in North American regions, the optimal Se consumption ranges from 52 to 64 $\mu\text{g}/\text{day}$ in Western Europe/France, and low levels of Se consumption range from 30 to 40 $\mu\text{g}/\text{day}$ in Eastern European countries/Poland (Zwolak, 2019). This metalloid is also known as a cancer chemopreventive compound and is indispensable for proper cell function (Zeng et al., 2005). Numerous mechanisms have been reported regarding the chemoprotective effects of Se, such as antioxidant protection, reduction in carcinogen metabolism effects, enhancement of the immune surveillance system, and inhibition of the angiogenesis process and cell cycle (Lu and Jiang, 2001; Zeng, 2009). Approximately 0.5–1 billion people worldwide suffer from Se deficiency (Jones et al., 2017), which makes them prone to several diseases, such as white muscle and Keshan disease (Shahid et al., 2018). Selenium deficiency occurs in humans when Se intake is < 40 $\mu\text{g}/\text{d}$ (Navarro-Alarcon and Cabrera-Vique, 2008; Winkel et al., 2011) and may cause reduced bone metabolism, growth obstruction, irregularities in thyroid function, reduced fertility, immune system complications and even cancer (Navarro-Alarcon and Cabrera-Vique, 2008; Gupta and Gupta, 2017; Chang et al., 2019). Inorganic Se is 40 times more toxic than organic Se (Vinceti et al., 2009). However, higher intake levels of Se > 400 $\mu\text{g}/\text{d}$ (Winkel et al., 2011) can lead to severe toxic effects in humans, such as skin lesions, nail and hair loss, nervous disorders, amyotrophic lateral sclerosis, diabetes, and paralytic symptoms leading to death (Fordyce, 2013; Chauhan et al., 2019).

3.3. Cadmium

Cadmium (Cd) is toxic. This nonessential transition element occurs directly below Zn in the periodic table due its chemical similarity with Zn; in compounds, Cd(II) exists as a divalent ion (Wuana and Okieimen, 2011; Soliman et al., 2019). This may be part of the reason for Cd toxicity; since Zn is a well-known essential element, the substitution of Cd for Zn can damage several functions of the metabolic process (Nzengue et al., 2011). Cadmium is present as an impurity in different products, such as detergents, phosphate-based fertilizers, and different petroleum-based products, and is also a component of acid rain, which increases soil acidification and thereby increases Cd mobility (Musa et al., 2017a).

Rice is a significant source of human Cd risk, and the basic chemistry of Cd in reducing soils is important (Shi et al., 2011). Both Cd and Zn primarily co-occur in sphalerite in robust solutions as CdS and ZnS (Leng et al., 2019). As soon as sphalerite is mixed in aerobic soils and sulfur oxidation by microorganisms occurs, soluble Cd and Zn are released into the ground and react with soils (Hooda, 2010). In flooded paddy soils, numerous chemical changes occur that may affect the phytoavailability of Cd and Zn (Pan, 2015). The reduction of the physicochemical contents of soil increases the concentration of the CdS complex. When flooded soils drain, a decrease in soil physicochemical contents and CdS complex concentrations occurs due to rapid oxidation and increased soil Cd phytoavailability; Zn has little effect on the yield of rice grains that contain a high level of Cd at harvest (Hooda, 2010). Rainfall during the rice grain filling period can slow paddy soil oxidation and reduce Cd accumulation in rice grains, causing variation in the Cd level in rice grains from year to year (Hu et al., 2013).

Excessive Cd intake may cause several human health effects, such as renal damage caused by proteinuria due to Cd severely affecting the enzymes that reabsorb proteins in kidney tubules (Matović et al., 2015). The most remarkable occurrence of Cd toxicity due to excessive dietary Cd intake happened in the Jinsu River Valley, Fuchu, Japan, where people were afflicted by itai-itai disease (Matović et al., 2015). Other health symptoms of Cd exposure include painful osteomalacia and kidney failure (Wuana and Okieimen, 2011).

3.4. Lead

Lead (Pb) is a bluish-gray metal that naturally occurs in soil and rocks; it belongs to the IV group in the periodic table with atomic number 82 (Scerri, 2019). Lead is usually found in metal form at concentrations ranging from 10 to 30 mg/kg in the Earth's crust. Pb is considered a chalcophile metal with a strong affinity for S as well as other elements, and is found in forms such as PbS, PbSO₄, PbCO₃, i.e., galena, lead sulfate, and lead carbonate, respectively (Wuana and Okieimen, 2011).

While only an insignificant amount of the total Pb existing in soil is bioavailable, the key pools of Pb in soils are in soil solution (Chen et al., 2009b). Lead adsorption on the surface of clay-humus particles can be found in compounds, precipitated forms, secondary iron (Fe) and manganese (Mn) oxides, basic earth carbonates, soil humus and silicate matrices (Hooda, 2010). Pb(II) complexes are mainly ionic, including Pb²⁺SO₄²⁻, and Pb(IV) complexes have a tendency to be covalent, such as tetra lead Pb(C₂H₅)₄; a few Pb(IV) compounds such as PbO₂ are considered active oxidants (Wuana and Okieimen, 2011).

Previous studies have confirmed that soil solution Pb is the primary source for plant uptake, with a balance occurring among the soil solution and the more 'exchangeable' or 'labile' pools of Pb in the soil medium (Hooda, 2010). In some soil solutions, Pb might be concentrated with calcium carbonate (CaCO₃) or phosphate (Huang et al., 2016). Elevated concentrations of Pb in the soil lead to the formation of pyromorphite (Pb₅Cl(PO₄)₃), which mainly occurs near the plant root system, and the solubility of Pb in soils primarily depends on pH (Hough, 2010). The increased pH in soil can precipitate Pb in forms such as carbonate, phosphate or hydroxide, and can also lead to the formation of moderately Pb-organic mixtures (Hough, 2010). Increasing soil acidity leads to increased solubility and mobilization of Pb, usually at a slower rate than the rate of Pb accumulation in the soil organic layer (Potysz et al., 2017).

Several studies have reported that Pb adsorption significantly depends on the types of ligands involved in the formation of hydroxyl compounds of Pb, such as (PbOH⁺) & Pb₄(OH₄)⁴⁺ (Hough, 2010; Farkas and Buglyó, 2017). However, it has been further suggested that the adsorption of these complexes on montmorillonite might be an interrupted cation exchange process, unlike that on kaolinite and illite (Hough, 2010). In soil solution, the higher sorption of Pb via manganese oxide, iron oxide, halloysite, and imogolite by kaolinite, montmorillonite, and humus was observed (Hough, 2010; Violante, 2013). High Pb intake can cause serious health effects, especially in children who are more vulnerable to Pb toxicity, leading to lower IQ, mental deterioration, impaired development, reduced attention span, and hyperactivity (Liu et al., 2013b). In adults, Pb exposure can cause memory loss, joint weakness, damage to the cardiovascular system, and other negative effects (Patrick and Toxicity, 2006).

3.5. Zinc

In soil systems, the mobility of Zn depends on the solubility of its speciated form and on the soil properties, including pH, cation exchange capacity, redox potential and the availability of metal species in the soil; under oxidizing conditions, zinc sulfide (ZnS) is the dominant species in soil (Kumpiene et al., 2008). However, ZnS is insoluble, and the mobility of Zn is low in oxidizing soils since, at higher soil pH levels,

Zn adsorption is reduced, resulting in the release of organic Zn complexes and increased concentrations of chelating agents in the soil system (Güngör and Bekbölet, 2010).

At high pH levels, the Zn in the soil solution is primarily precipitated as zinc hydroxide Zn(OH)₂ and/or zinc carbonate (ZnCO₃) (Prasad et al., 2016). In soil solution, clay and metal oxides have the capacity to sorb Zn and tend to retard its mobility (Zalewska, 2012); at pH 4, Zn is more mobile than that at pH 6.5 due to sorption (Güngör and Bekbölet, 2010). Goethite, iron hydroxide Fe(OH)₃, may also reduce Zn mobility in soil and increase retardation; this effect has also been reported with humic acid (Loganathan et al., 2012). Zinc can disturb the activity of soil organisms, including decreasing microbe and earthworm activity, which delays the breakdown of organic matter (Prasanna, 2017). Excessive Zn intake can cause acute and chronic toxic effects on human health, such as nausea, loss of appetite, vomiting, abdominal cramps, headaches, and diarrhoea (Deshpande et al., 2013).

3.6. Manganese

The most significant, dominant, and stable forms of Mn primarily persist in rock-forming minerals (He et al., 2010). The Mn²⁺ cation is considered a substitute for Fe²⁺ and Mg²⁺ in silicates and oxides (He et al., 2010). In the soil development process, Mn primarily accumulates as oxides and hydroxides; consequently, the total Mn concentration in agricultural soil is substantially higher than that of other elements (Shaheen et al., 2013). Mn-containing vital minerals are mostly ferromagnesian silicate minerals such as chlorite, biotite, and hornblende (Sarbas, 2013). During the mineral weathering process, Mn complexes oxidize under atmospheric environments, leading to the rapid release and transformation of Mn into the oxides or hydroxides of secondary minerals that accumulate in the soil system (He et al., 2010). Manganese oxides have an affinity to coprecipitate with Fe oxides (Tan et al., 2010); therefore, Mn oxides accumulate in soil horizons with Fe oxides and hydroxides (Vodyanitskii, 2009). Low concentrations of total Mn in soils persist in dissolved forms, primarily adsorbed on mineral surfaces or chelated with organic matter, and these forms contribute to the Mn pool that is available to plants (He et al., 2010). Mn (II) is the most dominant species of Mn in equilibrated soil solutions on exchange sites on soil surfaces (Essington, 2015). Mn(III) and Mn(IV) are mainly associated with solid soil phases and exist as oxides such as carbonates and phosphates (He et al., 2010). Mn²⁺, the primary plant-available form of Mn, is complexed with organic or inorganic ligands, and its concentrations in the soil system decrease with increasing pH (Rengel, 2000). Under anaerobic conditions, the Mn²⁺ concentration in soil solutions of extremely weathered soils, including oxisols and ultisols, might be a significant cause of agricultural crop toxicity, such as that in rice plant systems (He et al., 2010; Antoniadis et al., 2019b). The main inorganic complexes and pairs of ions, such as MnCO₃, MnHPO₄, MnSO₄, Mn(OH)⁺, and MnMnCl₂, in Mn-rich soil solution are readily soluble in the water system and are the most bioavailable (Hooda, 2010). In soil solution, Mn(II) forms complexes with organic materials, including humic acid, organic acid, and amino acid (Zhang et al., 2019).

However, the Mn²⁺ that is firmly bound to fulvic acid remains thoroughly hydrated, and the complex formed between Mn²⁺ and humic substances might be described as an outer-sphere complex where ligands do not directly bond with Mn electrons (He et al., 2010). Mn-organic ligand complexes can be quite labile and exist in crop plants and microbes (McCann, 2012). Although Mn plays a vital role in several physiological and biochemical processes in plants and humans, Mn deficiency can severely inhibit plant crop growth and human health (Soetan et al., 2010). Excessive Mn concentrations are toxic to plants and humans (Menezes-Filho et al., 2009). In humans, Mn toxic effects usually occur in the brain and respiratory tract, and significant symptoms of Mn toxicity include Parkinson's disease, hallucinations, forgetfulness, nerve damage, lung embolisms, and bronchitis (Pahade and

Sharma, 2015; Ferrante, 2013).

3.7. Nickel

In soil solution, nickel (Ni) occurs as the aqueous nickelous ion Ni^{2+} , which can form complexes with organic/inorganic ligands or with suspended mineral colloids (Ma and Hooda, 2010). Organic complexes, rather than different inorganic species such as Ni^{2+} , NiCO_3 , H_2CO_3^+ , and NiOH^+ , are typically the most dominant Ni species in soil solution (Ma and Hooda, 2010).

The behaviour of Ni is also pH-dependent; at low pH, Ni occurs in the form of Ni^{2+} , whereas in neutral to high-pH soil solutions Ni precipitates as stable nickelous hydroxide $\text{Ni}(\text{OH})_2$ compounds (Wuana and Okieimen, 2011; Musa et al., 2017b). Nickel is radially dissolved in acidic soil solutions to form Ni^{3+} , and very in basic soil solutions it forms nickelate compounds (HNiO_2) which are soluble in water (Wuana and Okieimen, 2011). Notably, the total Ni content in soils, the significant sources of Ni in soils, including geogenic and anthropogenic sources, and the physicochemical characteristics of soils influence the partitioning of Ni between solution and its solid phase in soil; these factors control Ni mobility and bioavailability (Ma and Hooda, 2010; Antoniadis et al., 2017). Nickel occurs in crucial minerals, mainly allied with oxides of Mn and Fe and layered silicates, such as chlorite, garnierite and vermiculite (Echevarria, 2018). In serpentine soils with a high concentration of Ni and basic pH, Ni can partially bind with carbonates (Vithanage et al., 2014). In solution, Ni has an affinity to rapidly adsorb on the surfaces offered by the soil solid phase such as layered silicates, Fe(Al), Mn-oxides, and organic matter (Violante et al., 2010), and can form inner or outer-sphere surface complexes (Ma and Hooda, 2010; Yang et al., 2011).

Nickel adsorption by the formation of outer-sphere surface complexes is radially exchangeable because Ni species occur as hydrates and loosely attach to outer-sphere surfaces of the solid soil phase through electrostatic force and by H-bonding between hydration shells and surface oxygen (Rinklebe and Shaheen, 2017). In contrast, Ni species adsorbed on the inner-sphere surface complexes are partly dehydrated or dehydrated species (Ma and Hooda, 2010).

Compared with other trace elements, the adsorption affinity of Ni^{2+} ions for soil mineral materials is usually lower than that of Zn^{2+} , Pb^{2+} , and Cu^{2+} (Igberase et al., 2018). The stoichiometric precipitates, i.e., nickel phosphate $\text{Ni}_3(\text{PO}_4)_2$ and nickel ferrite (NiFe_2O_4), are seldom responsible for controlling the Ni concentration in soil solution, even in severely fertilized and contaminated soils (Ma and Hooda, 2010). However, Ni precipitates are mostly detectable on clay mineral surfaces and Al oxides at sorption densities lower than the theoretical monolayer of Ni sorption (Abbas et al., 2014). Moreover, these precipitates can form at circumneutral pH with saturated concentrations of Ni in solution via the solid phases of Ni (Konhauser et al., 2009).

The most important factor affecting the behaviour of Ni in soils appears to be pH, though several characteristics, including the clay content and Fe and Mn oxides in soils, are considered of secondary importance (Du Laing et al., 2009). The mobility of Ni in the soil system increases as the pH and cation exchange capacity decrease (Ehsan et al., 2014). Several other factors, including waterlogging (redox), organic composts, and sewage sludge in the soil, also affect the fate and transport behaviour of Ni in soils (Adriano, 2013). At low levels, Ni is considered an essential element for plants, humans, and animals, but a high intake of Ni leads to serious human and animal health effects and is toxic to plants (Jaishankar et al., 2014). The most significant human health problems caused by Ni exposure include chronic bronchitis, lung cancer, nasal sinus damage, and reduced lung function (Salnikow and Zhitkovich, 2007).

3.8. Chromium

Chromium (Cr), a transition metal, does not exist naturally in

elemental form, only in compounds (Scerri, 2019). The significant natural source of Cr in soil is primarily ore products such as mineral chromite, while anthropogenic sources of Cr include electroplating and Cr-containing industrial waste (Saha et al., 2011). Cr(VI) is the main form of Cr found at contaminated sites. Its presence depends on the pH and redox conditions. Cr also occurs in the (+III) oxidation state since Cr(IV) is significantly dominant in shallow aquifers where oxidizing conditions exist (Guertin et al., 2016). The main Cr species, such as chromate (CrO_4^{2-}) and dichromate ($\text{Cr}_2\text{O}_7^{2-}$), can be precipitated in the presence of metal cations, including silver, barium, and lead ions (Ag^+ , Ba^{2+} & Pb^{2+} respectively) (Wuana and Okieimen, 2011).

CrO_4^{2-} and $\text{Cr}_2\text{O}_7^{2-}$ adsorb on soil surfaces, particularly on aluminium and iron oxides (Dotaniya et al., 2014). At low pH < 4, Cr(III) is the more dominant form of Cr, and Cr(+III) may form complexes with ammonia, hydroxide, chloride, fluoride, cyanide, or sulfate (NH_3 , OH^- , Cl^- , CN^- or SO_4^{2-} , respectively) and with soluble organic ligands (Wuana and Okieimen, 2011). Cr(VI) is considered a more toxic and more mobile form of Cr. The mobility of Cr(III) is also decreased through adsorption on clay and oxide minerals at low pH < 5, and it is less soluble at pH > 5 due to the formation of the chromium hydroxide ($\text{Cr}(\text{OH})_3$)_(s) complex (Wuana and Okieimen, 2011; Jiang et al., 2018).

The mobility of Cr depends on the sorption features of the soil, the clay content in soil, the iron oxide content, and the availability of organic matter (Dotaniya et al., 2014). The soluble and precipitated forms of Cr are transported by surface runoff to surface water bodies (Nordstrom, 2011), while soluble and unadsorbed Cr complexes can leach from the soil to aquifers; the leaching ability of Cr(VI) increases as the soil pH increases (Wuana and Okieimen, 2011). Human health risks associated with elevated Cr intake include respiratory distress, irritation, loss of body weight, and allergic dermatitis (Teklay, 2016).

4. Mechanisms of trace element uptake, translocation, and accumulation in paddy soil-rice systems

A range of different trace elements generally exists in soil systems. Paddy plants can acquire the required trace elements from the soil as micronutrients (ar Hashmi and Varma, 2018). The uptake, translocation, and accumulation of different trace metals in different paddy plant parts followed the order root > shoot > leaf > husk > grain, and the concentrations of these trace metals in plant parts depend on the exposure level (Khanam et al., 2019). Trace metals such as Zn and Cu are essential for paddy plant growth and sustenance; however, higher uptake, translocation, and accumulation of essential or nonessential trace metals can lead to toxicity and inhibit the growth and production of paddy rice (Kwon et al., 2017). The root zone (rhizosphere) of paddy plants and soil microbes play a significant role in trace metal availability as well as uptake and translocation to paddy plant systems (Dotaniya and Meena, 2015). Trace metals can be bound to soil particles or precipitated, which might make a crucial portion of trace metals insoluble or unavailable to paddy plants (Dotaniya and Meena, 2015). Several microbial communities in the rhizosphere of paddy plants can catalyse redox reactions in the soil system that can change the mobility of trace metals and the tendency of paddy plants to take up metals (Zhang et al., 2015). In metal-rich soils, after the biodegradation or biotransformation of trace metals, plants can accumulate specific trace metals in their tissues (accumulators) or, in the same way, restrict metal uptake into their biomass (excluders) (Gupta et al., 2014; Moreno-Jiménez et al., 2012). However, paddy plants are considered strong trace metal accumulators (Greger, 2008).

In paddy plants, the uptake, translocation, and accumulation of essential, toxic and potentially toxic trace elements in their biomass depends on environmental and biogeochemical factors such as the availability of soil microbes; the availability of trace elements, exudates, and chelating agents produced by plants; the pH changes induced by oxidation-reduction reactions; and the availability of transporters and cotransporters embedded in the plant cell plasma membrane

(Gupta et al., 2014; Verkleij et al., 2009). Specific transporter proteins generally involved in trace metal uptake and translocation, such as adenosine triphosphate proteins (ATPases), produce electrochemical gradients and consume energy (Gupta et al., 2014; Tangahu et al., 2011).

The co- and anti-transporter proteins generally use the electrochemical gradients generated by ATPases, and different ion channel proteins facilitate the normal transport of ions in plant cell systems (Gupta et al., 2014, 2016). Nevertheless, several studies have reported that the co-transportation of cations is a common phenomenon, and potentially toxic trace metals that do not have any known biological or beneficial function, such as As, Cd, and Pb, are mostly transported throughout paddy systems via transportation systems (Gupta et al., 2014). In the root zone of paddy plants, the immobilization of and controlled access to trace metals might be achieved through root exudates (UdDin et al., 2015) that chelate trace metals to prevent their uptake into root cells, such as Ni chelation; exudates can also bind Cu or Zn in the apoplast (Gupta et al., 2014). Therefore, the chemical properties of the rhizosphere and the cell walls in the roots may control trace metal uptake, translocation, and accumulation in paddy plants (Hu et al., 2011). Transporter proteins in the plasma membrane of plant root cells and the tonoplast of the cytoplasm (Weston et al., 2012) can maintain the physicochemical concentrations of and stress responses to trace metals, contributing significantly to trace metal homeostasis in the plant (Gupta et al., 2014). In paddy plant systems, approximately 13 putative ZIPs are a significant group of trace metal transporters that are primarily involved in the translocation of divalent cations across plant cell membranes (Lee et al., 2010). ZIPs also exist in other plant species as well as in microbes, fungi, and animals (Komal et al., 2015).

These specific proteins are topologically analogous to the eight transmembrane protein (TP) domains, with the N- and C-termini uncovered to the apoplast, and a histidine-rich cytoplasmic circle skirted via transmembrane domains (III) and (IV) is commonly involved with the binding of trace metals (Gupta et al., 2014; Milner et al., 2013). In

paddy plant systems, As uptake depends on the As concentration gradient between the source and sink (Abbas et al., 2018). The As uptake in the plant cell (Fig. 2a) mainly depends on As species such as arsenate (As(V)) and uses different phosphate transporters (Pi) that belong to the PHT1 family because P is chemically similar to As(V) (Table S2) (Moreno-Jiménez et al., 2012).

Arsenite (As(III)) uses silicon (Si) transporters such as Lsi1 (aquaporin NIP2;1) and Lsi2 (an efflux carrier) due to the resemblance between As(III) and Si (Bastías and Beldarrain, 2016). As(III) is absorbed by the aquaglycoprotein nodulin-like natural proteins (NIPs) (Bastías and Beldarrain, 2016). Under Si deficiency, the expression of Si influx transporters (Lsi1 & Lsi2) increases (Ma and Yamaji, 2008). The accumulation of Si in plant cells is mostly controlled by the Lsi1 and Lsi2 transporters, which are mostly located at the proximal or distal flanks of epidermal and endodermal cells; these cells help in the transportation of As across the plant's cells and tissues (Abbas et al., 2018). Several studies have suggested that the reduction of As occurs mainly in root cells before its transport to xylem and the rest of the plant (Zhao et al., 2009a). The main As species, As(III) and As(V), are primarily found in the xylem sap of plants (Finnegan and Chen, 2012). Of the low concentration of total As absorbed through plant roots, only a very small amount is sequestered in leaf, shoot, and grain vacuoles due to As reduction, and the sequestration mechanisms are similar to those of the roots (Bastías and Beldarrain, 2016). Hence, the presence of As(III) and As(V) in phloem is a requirement for its distribution to other parts of the plant (Chen et al., 2017), while Se translocation in plant shoots, leaves, and grains depends on the rate of transpiration and the rate of xylem loading (Gupta and Gupta, 2017; Renkema et al., 2012).

Selenium uptake in the plant system is facilitated by transporters (Fig. 2b). Selenite (Se(IV)) and selenate (Se(VI)) are transported through phosphate and sulfate channels (Shahid et al., 2018). Se(VI) enters the plasma membrane of plant root cells using sulfate transporters (Lin et al., 2012), and Se(IV) and P compete to cross the plasma membrane of paddy plant root cells using P transporters (Winkel et al.,




Fig. 2. The uptake, translocation, and accumulation mechanism of trace elements, (a) arsenic, (b) selenium, (c) cadmium, (d) lead, (e) zinc, (f) manganese (g) nickel, and (h) chromium in paddy soil-rice system.

Table 2
Reported trace element concentrations ranges minimum-maximum (mean) mg/kg in grain in different countries.

Country	Locations	Cd	Cu	Ni	Pb	Zn	Cr	Mn	Se	As	References
China	Wenling	0.1–0.5 (0.1)	0.7–5.8 (3.1)	0.05–1.7 (0.2)	-	11.4–35.4	-	-	-	-	Zhao et al. (2010)
	Shengzhou	0.01–0.2 (0.1)	1.3–4.7 (3.2)	0.1–1.5 (0.4)	0.05–0.2 (0.1)	18.5–27.8 (23.5)	0.1–1.4 (0.2)	-	-	-	Zhao et al. (2009b)
	Zhejiang	0.1–0.112 (0.04)	-	-	0.01–0.2 (0.1)	9.45–22.0 (14.3)	-	-	-	0.1–0.3 (0.1)	Huang et al. (2013)
	Nanxun	0.01–0.10 (0.01)	1.1–4.2 (2.5)	0.02–1.4 (0.1)	-	-	-	-	-	-	Zhao et al. (2015b)
Hunan	Nanhu	0.01–6.2 (0.6)	-	-	1.2–4.8 (0.05)	21.1–27.1 (23.2)	0.2–0.4 (0.2)	-	-	0.01–1.0 (0.1)	Lu et al. (2018)
	Tongxiang	-	2.9–8.9 (5.5)	-	0.1–0.5 (0.2)	18.1–27.5 (22.6)	0.3–0.3 (0.3)	-	-	-	Zeng et al. (2011)
Xiaoshan		-	2.4–5.5 (4.1)	-	0.2–0.6 (0.3)	17.7–25.2 (21.3)	0.1–0.3 (0.2)	-	-	-	-
		-	2.8–6.7 (4.7)	-	0.2–0.5 (0.3)	-	-	-	-	-	-
Shimen		0.01–0.7 (0.3)	-	0.3–1.2 (0.7)	0.01–0.03 (0.02)	-	0.03–0.11 (0.1)	-	-	0.3–1.2 (0.5)	Zeng et al. (2015)
	Fenghuang	0.01–2.1 (0.3)	-	0.1–1.4 (0.5)	0.003–0.02 (0.01)	-	0.05–0.14 (0.1)	-	-	0.2–0.4 (0.3)	-
Xiangtan		0.01–0.691 (0.3)	-	0.1–0.8 (0.5)	0.01–0.10 (0.04)	-	0.1–0.5 (0.2)	-	-	0.1–0.3 (0.2)	-
	Kaiyang	-	-	-	-	-	-	-	0.04–0.14 (0.08)	-	Zhang et al. (2014)
Yangtze River		-	-	-	-	-	-	-	0.02–0.04 (0.03)	-	Cao et al. (2001)
	Enshi	-	-	-	-	-	-	-	0.08–9.7 (2.73)	-	Sun et al. (2010)
Wanshan		-	-	-	-	-	-	-	0.02–0.7 (0.1)	-	Zhang et al. (2012a)
	Rugao	0.1–0.25 (0.1)	0.3–2.2 (1.1)	-	0.24	-	0.03	0.016–0.40 (0.09)	-	-	Sun et al. (2009)
Narora		0.9–5.7 (1.9)	0.04–0.41 (0.4)	0.06–1.0 (0.6)	5.5–20.2 (11.6)	0.1–0.9 (0.6)	-	-	-	-	Singh et al. (2014)
	Fatehabad	-	2.1–4.6	0.13–1.1	29–1529 (173)	3.21–10.3	-	-	-	-	Yadav et al. (2017)
West Bengal	Market	9.4–178.0 (27.5)	-	-	-	-	-	-	-	-	Roychowdhury et al. (2003)
	purchased	-	-	-	-	-	-	-	-	-	Rahman et al. (2014)
Bengal delta		-	-	-	-	-	-	-	-	0.07–0.31	Stroud et al. (2011)
	Punjab	-	-	-	-	-	-	-	4.83–8.0	-	Khanam et al. (2019)
Bangladesh	Market	69–77 (73.0)	1.3–1.9 (1.6)	76–163 (105)	5–28 (19.0)	12.1–14.9 (13.4)	41–217 (119)	13.2–16.0 (14.7)	-	-	Rahman et al. (2014)
	purchased	(0.1)	-	-	(0.3)	-	-	-	-	0.41–0.98	Khanam et al. (2019)
Market	purchased	-	-	-	(3.19)	-	-	-	-	0.02–0.33	Khanam et al. (2019)
	Market	-	-	-	(0.7)	-	-	-	0.07–0.16	-	Khanam et al. (2019)
Thailand	Market purchase	8.2–16.0 (13.4)	-	-	23.1–68.1 (37.0)	-	-	-	0.05–0.10	-	Shraim (2017)
	Market purchase	0.1–8.9 (4.5)	1.9–3.6 (2.6)	77–180 (118)	3–168 (67.0)	14.5–17.1 (16.2)	48–113 (79.0)	7.3–8.7 (7.8)	0.05–0.25	(0.15)	Rahman et al. (2014) and Shraim (2017)
Pakistan	Shakargath	0.2–1.6 (0.9)	-	38.6–52.6 (46.3)	38.6–55.9 (45.7)	-	4.6–12.2 (7.0)	-	-	-	Tariq and Rashid (2012)
	Market purchase	8.7–17.1 (7.5)	1.0–9.4 (2.9)	61–356 (166)	16–1248 (375)	10.9–24.5 (17.1)	15–465 (144.0)	9.2–51.7 (24.4)	-	-	Rahman et al. (2014)
Malaysia	Koita	0.03–0.18	0.3–0.4	-	-	0.4–0.7	0.2–1.3	-	-	0.10–0.77	Yap et al. (2009)
	Four mining areas	0.01–0.98 (0.17)	2.0–29.6 (4.7)	-	0.01–3.34 (0.8)	7.3–38.0 (16.8)	-	-	-	(0.25)	Kwon et al. (2017)
Iran	Several locations	(0.64)	(1.1)	(0.776)	(0.64)	(28.6)	(0.4)	-	-	(0.05)	Jafari et al. (2018)
	Ghaemshahr	-	-	(1.0)	(6.1)	-	(2.4)	-	-	(0.94)	Asgari et al. (2009)
USA	Market	(0.02)	-	-	0.01–0.09	-	-	-	(0.02)	0.2–0.7	Khanam et al. (2019)
	purchased	-	-	-	-	-	-	-	-	-	Khanam et al. (2019) and Zhang et al. (2014)

(continued on next page)

Table 2 (continued)

Country	Locations	Cd	Cu	Ni	Pb	Zn	Cr	Mn	Se	As	References
Japan	Market purchased Aomori	(0.05) 0.004–0.1 (0.03)	- 0.6–3.9 (2.1)	- 0.02–0.4 (0.11)	(0.002) 0.001–0.09 (0.01)	- 13.0–19.0 (16.0)	- 0.001–0.05 (0.01)	- 5.4–18.0 (9.7)	0.03–0.109 0.001–0.006 (0.003)	(1.8) 0.05–0.15 (0.09)	Khanam et al. (2019) Tsukada et al. (2007)

2015). Se^{IV} and P compete and enter the plasma membrane of plant root cells by P transporters (Winkel et al., 2015). The P-induced increase in Se bioavailability most likely occurs through the exchange of Se at sorption sites, therefore increasing Se mobility and uptake in the plant (Shahid et al., 2018). In plant cells, Se typically accumulates in vacuoles and effluxes with S transporters present in the tonoplast (Mazaj et al., 2008; Hawkesford and De Kok, 2006). Selenium, after entering plant cells with the help of the sulfate transporter, is translocated to other parts of the plant, i.e., the shoots, leaves, and grain (Bitterli et al., 2010), and is metabolized in plastids through the S integration pathway into selenomethionine (SeMet) or selenocysteine (SeCys); the S chemical analogue with Se can become more methylated and evaporate in a nontoxic form (Pilon-Smits and Quinn, 2010).

In paddy plants, the uptake, translocation, and accumulation of potentially toxic trace metals such as Cd from the paddy soil-rice (root cells to shoots) system are generally facilitated by ZIP (OsIRT1) transporters (Uraguchi and Fujiwara, 2012). These ZIP transporters are essential for paddy plants, and the plant capacity for critical trace metal accumulation is associated with the level of ZIP expression (Lee and An, 2009). In the Cd uptake, translocation, and accumulation mechanism in the paddy soil-rice system, Cd travels from the soil to the above-ground parts of the rice plants. Cadmium is mainly absorbed through rice root cells from the paddy soil and is translocated to other rice parts (shoots and leaves), finally accumulating in the rice grains (Fig. 2c). In the Cd absorption process, xylem plays a significant role in the transportation of Cd from root cells to shoot cells, and the phloem pathway is used in the translocation of Cd to the rice grains (Khanam et al., 2019). ZIP transporter-like OsIRT1 and OsIRT2 might show influx activity for Cd²⁺ ions. Therefore, it has been reported that OsIRT transporters play a significant role in Cd uptake through roots in paddy fields (Khanam et al., 2019). However, NRAMP3 and NRAMP4 transporters and their cotransporters also play a crucial role in Cd transport and homeostasis as well as in the resistance to the toxic effects of Cd in paddy plants (Kim and Guerinot, 2007). Furthermore, the high concentrations of Cd and Mn in paddy plants are primarily facilitated by NRAMP5 transporters (Sasaki et al., 2012).

Additionally, the Fe transporter-like OsNramp1 also functions as Cd influx transporters in the plant plasma membrane. It has been reported that Cd is readily transported in paddy plants from root cells to shoot cells through culms and slants and accumulates in nodes (Fujimaki et al., 2010). The ABC transporter family is involved in Pb and Cd transportation from outside the plasma membrane to inside the plasma membrane (Uraguchi and Fujiwara, 2012; Song et al., 2014). Pb²⁺ ions usually occur in paddy soil solution and can enter paddy plants through the translocating water uptake (Fig. 2d). In paddy plants, root cell H⁺ + ATPase and H⁺ + drivers can directly take up Pb.

Furthermore, the transpiration process also plays a significant role in pulling or pushing Pb²⁺ from root cells to shoots through the xylem and from the shoots to leaf cells through vascular flow; Pb translocation to paddy rice grains is still not well defined (Khanam et al., 2019). ZIP family transporters such as IRT1, ZRT1, and ZRT2 facilitate the uptake, translocation, and accumulation of different cations, such as Zn²⁺, Mn²⁺, Cd²⁺, & Ni²⁺, in paddy systems (Fig. 2 e, f & g) (Dutta et al., 2018; Mani and Sankaranarayanan, 2018). Furthermore, the Arabidopsis proteins ZIP1 and ZIP2 transporter-like (AtZIP1 & AtZIP2) are generally involved in the translocation of both Mn and Zn from the root to the shoot. They may also play a significant role in the process of remobilization and distribution at the cellular level (Ahmadi, 2017; Pinto et al., 2014).

Another family of trace metal transporter-like natural resistance-associated macrophage proteins (NRAMPs) is expressed in paddy plant roots and shoots, and these transporters are involved in the transport of several trace metals, such as Ni²⁺, Mn²⁺, Zn²⁺, Cd²⁺, Cu²⁺, & Co²⁺, across paddy plant cell membranes (Gupta et al., 2014; Tiwari et al., 2014; Ishimaru et al., 2012). However, given the availability of Cr and its species, such as Cr(III) and Cr(VI), in the paddy soil system, paddy

plants have an affinity for translocating and accumulating both, while the mechanism of Cr uptake in the paddy soil-rice system is still unclear. Generally, Cr is taken up by plants *via* specific transporters for the absorption of the essential trace metal ions that are required for plant metabolic activity.

Cr(III) uptake in plant systems occurs through passive mechanisms, and Cr(VI) uptake occurs through active processes, usually *via* phosphate or sulfate transporters (Fig. 2h), due to the structural similarity of Cr(VI) to phosphate and sulfate (Shahid et al., 2017a). However, the uptake, translocation and accumulation mechanisms of trace metals such as U, Tl, Rb, and Sr are still unclear.

5. Status of trace element contamination in paddy soil-rice systems and risks to rice-growing countries

Paddy crops show substantial tolerance of trace element toxicity; however, trace metal accumulation in different plant parts, *i.e.*, the roots, shoots, leaves, and grains of paddy plants, in varying concentrations has been reported (Clemens and Ma, 2016; Hseu et al., 2010). From this perspective, field investigations are particularly significant, as the findings of such studies could be efficiently used to measure the status of trace element contamination and accumulation in paddy systems (Arunakumara et al., 2013).

China is a leading rice-growing country in the world. Trace elements resulting from various human activities contaminate approximately 20 million h in China, which covers 20 % of the total agricultural area. Paddy cultivation is concentrated in southern China, which also has a highly developed economy and dispersed common nonferrous trace metals, which may seep with nonferrous trace metals into paddy soil systems (Arunakumara et al., 2013). Thus, trace metal contamination in paddy soil-rice systems has become a severe concern in China (Zhao et al., 2010).

A study was conducted on paddy soils from Fenghuang, Shimen, and Xiangtan in Hunan Province, China. The mean value of Cd in the paddy soils was five times higher than the WHO permissible limits for Cd (0.35 mg/kg), whereas the maximum Pb, Ni, and As (48.7, 262 & 45.2 mg/kg) concentrations were higher than the permissible limits (35, 50, & 10, respectively) (Table 2). Paddy soil pollution was mainly associated with anthropogenic activities such as fertilizer and pesticide application during agricultural production. Consistent with the paddy soil results, the mean value of Cd in rice grains from Fenghuang, Shimen, and Xiangtan (0.28, 0.32, & 0.34 mg/kg, respectively) (Table 3) exceeded the national permissible limits (0.2 mg/kg), whereas the levels of As, Ni, Pb, and Mn contamination in rice grains with the average values of the transfer factors (TFs) for Cd, Cr, As, Ni, Pb, & Mn decreased in the order of Cd > Mn > As > Ni > Cr > Pb. Cd had the highest TF of 0.55 in rice grains, confirming the 81.2 % increase in cancer risk in the population of Hunan Province, China, which is mainly attributed to the consumption of Cd-contaminated rice (Zeng et al., 2015).

Recently, a similar study conducted in the industrial and urban areas of Hunan Province reported that the mean concentrations of Cd, Pb, and As in paddy soil systems were 1.08, 52.8, and 19.5 mg/kg, respectively, and 68.9 % of rice grain samples exceeded the permissible limit for Cd concentration in rice grains (Fig. 3). The maximum TFs of Pb, As, and Cd from paddy soil to rice grains were reported as 0.023, 0.044, & 4.13, respectively, and the cancer risk of the studied metals in rice grains followed the order Cd > As > Pb; these metals pose a serious health risk to the population (Lu et al., 2018). In a study conducted by Zeng et al. (2011), paddy soil and rice samples collected from three major rice-growing areas (Tongxiang, Nanhu, & Xiaoshan) in Zhejiang Province, China, reported that significant correlations between the trace metal contents in paddy soils and rice plants were only observed in Mn and Cu in rice shoots and Zn and Mn in rice grains, concluding that the trace metal concentrations in the paddy soil-rice system were not acceptable (Zeng et al., 2011).

In Zhengzhou City, China, a field experiment conducted by Liu et al.

(2007) evaluated the uptake and translocation of As, Cd, Cr, and Pb in parts of paddy plants and found the accumulation of trace metals in rice grains to be comparatively lower than that in roots and shoots. They confirmed that the levels of As, Cd, Cr, and Pb in the edible rice grains were much lower than the Chinese national limits for food; therefore, no threat to human health in terms of trace metal toxicity was involved with consuming the rice (Liu et al., 2007).

In contrast with this study, Luo et al. (2012) studied the effect of agronomic practices on trace metals in soils and vegetation mainly in the water conservation area along the East Dongjiang River, South China, and found that Cd and Pb concentrations in rice grain were much higher than the Chinese national limits for food; hence, they identified a potential human health risk mainly associated with the long-term consumption of contaminated rice (Luo et al., 2012). According to their study, which covered an area of approximately 13,600 km² in Heyuan, Huizhou, Guangzhou, and Dongguan, they further suggested that the average Cd concentrations in paddy soils also exceeded the Chinese national maximum permissible limits (Arunakumara et al., 2013).

Paddy soils adjacent to a coal mine in Guizhou Province, China, were found to contain higher concentrations of As (83 mg/kg). In rice plants in the same area, As concentrations ranged from 6.3 to 44.7 mg/kg in the roots, from 2.2 to 8.9 mg/kg in the shoots, and from 0.3 to 1.7 mg/kg in rice grains, with an average As concentration of 0.8 mg/kg (Zhang et al., 2013). The distribution and translocation of Se in a paddy soil-rice system was assessed by Sun et al. (2010) in Enshi, Hubei Province, which is typical of Se-rich areas in China. The reported concentration of Se in paddy soils ranged from 0.5 to 47.7 mg/kg, whereas the Se concentration in the rice bran was 1.94 times higher than that in the polished rice grains. The total Se concentrations in the paddy plant parts were found in the order straw > bran > whole-grain > polished rice > husk and exceeded the Chinese national maximum permissible limits for Se in rice (0.3 mg/kg) (Sun et al., 2010). In contrast, in paddy fields near an old in Tan Long, Vietnam, Dang et al. (2008) reported high concentrations of trace elements, such as Cd, ranging from 2.3 to 42.9 mg/kg, and Pb, ranging from 1271 to 3953 mg/kg. In Chi Dao, Vietnam, paddy fields situated less than 20 m away from smelters were found to have total Pb and Cd contents ranging from 2000 to 10,000 and 2.5 to 56.5 mg/kg, respectively (Chu, 2011).

Due to the increased population and industrialization, Japanese paddy fields suffer from severe paddy soil contamination by trace metals (Arunakumara et al., 2013;). In Japan, metal contamination in paddy soils is primarily attributed to wastewater from mines that is used for irrigation purposes and uncontrolled nonferrous emissions from metal refining plants (Makino, 2007). According to Makino et al. (2010), the total concentrations of trace metals such as Zn, Cr, Ni, Cd, and Pb (99.0, .64.0, 39.0, 0.45, & 29.0 mg/kg, respectively) in the paddy soil system were significantly elevated compared with the natural concentrations (89.0, 58.0, 26.0, 0.3 & 24.0 mg/kg, respectively). Consequently, an elevated level of Cd in the paddy soil-rice system has been frequently reported, and it is recognized that rice is a significant source of human Cd intake in Japan (Arao et al., 2010; Uruguchi and Fujiwara, 2013).

The Cd concentration in Japanese rice has been found to range from 0.02 to 1.06 mg/kg (Nogawa et al., 2004). The average nutritional consumption of Cd by the Japanese population is approximately 3.0 µg Cd/kg body weight/week, which exceeds the acceptable weekly consumption (2.5 µg Cd/kg body weight/week) defined by the European Food Safety Authority (EFSA) (Uruguchi and Fujiwara, 2012).

A study conducted by Tsukada et al. (2007) in Aomori Prefecture, Japan, reported the status of trace element contamination in a paddy soil-rice pf. He used the paddy soil-to-paddy plant TF and suggested that the TFs for all investigated trace elements, excluding Pb, were within two orders of magnitude. According to Yap et al. (2009), a study conducted in Kota Marudu, Malaysia reported that trace metals such as Cd, Cr, Mn, Zn, Cu, & Pb can accumulate in different parts of the paddy

Table 3

Trace element content and translocation factor (TF) from paddy soil to different parts (roots, shoots, & grains) of the rice plant.

Metal	Metal level in soil mg/kg	Metal level in root mg/kg	Metal level in shoot mg/kg	Metal level in grains mg/kg	Metals translocation factor (TF) from paddy soil to rice plants parts				References
					Soil to root	Root to shoot	Shoot to grains	Soil to grain	
As	5.42	22.77	0.68	0.06	4.2	0.03	0.09	0.011	Singh et al. (2011b)
Cd	0.86	6.16	0.62	0.01	7.19	0.1	0.02	0.012	
Cu	23.22	2.09	0.17	0.05	0.09	0.08	0.29	0.002	
Pb	18.65	7.09	1.84	0.52	0.38	0.26	0.28	0.028	
Cd	2.39	16.32	3.35	0.85	6.83	0.21	0.25	0.356	Kibria et al. (2012)
Cu	59.13	32.67	5.52	2.79	0.55	0.17	0.51	0.047	
Pb	40.91	29.46	4.45	0.84	0.72	0.15	0.19	0.021	
Zn	305.03	80.75	27.58	2.3	0.26	0.34	0.08	0.008	
Cd	35.02	0.85	0.85	0.85	–	–	1.0	0.024	Tariq and Rashid (2012)
Pb	1065	–	48.81	45.75	–	–	0.94	0.043	
Cr	37.8	–	9.17	6.99	–	–	0.76	0.185	
Co	747.2	–	71.58	49.89	–	–	0.7	0.067	
Ni	85.84	–	54.5	46.34	–	–	0.85	0.539	
Cd	0.58	–	1.59	0.48	–	–	0.29	0.001	Luo et al. (2012)
Cu	20.4	–	0.3	0.24	–	–	0.8	0.012	
Pb	27.9	–	0.07	0.02	–	–	0.29	0.001	
Ni	20.4	–	0.08	0.09	–	–	1.13	0.004	
Cd	0.17	–	–	0.01	–	–	–	0.059	
Pb	29.6	–	–	0.05	–	–	–	0.002	Cao et al. (2010)
Se	11.46	11.82	1.35	1.88	0.96	0.18	0.71	0.12	Chang et al. (2019)
Cd	4.64	–	–	1.81	–	–	–	2.5	Liu et al. (2016)

plant, furthermore suggesting that the concentrations of investigated trace elements in rice grains were below the maximum permissible limits specified by the Malaysian Food Act and Food Regulations.

In western Bangkok, Thailand, a study conducted by Chinoim and Sinbuathong (2010) examined the status of trace metal (Cr, Cd, Ni, Pb, Zn, & Cu) contamination in paddy soils and reported that the concentrations of the studied trace metals were much lower than the acceptable limits. In contrast, Shraim (2017) studied the status of trace metals in market-purchased Thai rice and reported Cd and Pb

concentrations that exceeded acceptable limits. A study conducted by Jo and Koh (2004) on approximately 4047 paddy soil samples collected from different locations in Korea reported that the status of trace metal contamination (Zn, Pb, Cd, & Cu) ranged from 0.1 to 96.7, 0.01 to 66.4, 0.1 to 1.01, and 0.1 to 41.6 mg/kg, respectively.

A recent study conducted by Kwon et al. (2017) in a paddy soil-rice system adjacent to four mining areas (Dalsung mine, Yeongdae mine, Sambo mine, and Munmyung mine) in Korea reported that the average concentrations of Cd, Cu, Pb, Zn, and As in paddy soils and grains were




Fig. 3. Trace element arsenic (As), selenium (Se), cadmium (Cd), lead (Pb), zinc (Zn) manganese (Mn), nickel (Ni), and chromium (Cr) contamination in grains of major rice-growing countries.

comparatively higher than the worldwide acceptable limits defined by numerous researchers (Tables 1 and 2). Furthermore, an acceptable daily intake (ADI) of As and Cd in rice grains of up to 50 % and 80 %, respectively, is recommended by the Joint Food Additive and Contaminants Committee and the World Health Organization (FAO & WHO), respectively. Consequently, the regular consumption of contaminated rice from these mining areas can lead to severe human health problems in the local population. In Iran, a review on the status of trace element contamination (As, Cr, Pb, Cd, & Ni) in rice grains and an assessment of the associated human health risks noted that the trace metal concentrations were highest in rice grains from northern areas of Iran, primarily Mazandaran & Gilan provinces (Abtahi et al., 2017). Furthermore, minimum and maximum levels for trace metals (As & Ni) were established in Iran, and the ranked order of trace metals by their total hazard quotient (THQ) in the human health risk assessment was $As > Cr > Pb > Cd > Ni$ (Abtahi et al., 2017).

In Bangladesh, a study conducted by Ahsan et al. (2009) stated the current scenario for contamination by As and other trace metals in the floodplain agricultural field of the Faridpur and Dhamrai study areas. This study reported that the average As concentration in the Faridpur soil system exceeded the WHO acceptable limits (10 mg/kg) by three times, whereas the average levels of other trace metals (Cr, Co, & Cu) in both study areas also exceeded the WHO acceptable limits (70, 8, & 30 mg/kg, respectively); however, the Se concentration was much lower than the WHO acceptable limit (0.7 mg/kg). Furthermore, in both study areas, the groundwater contaminated with As and other trace metals also led to elevated concentrations of these trace metals in soils because of the predominant use of irrigation with the polluted groundwater (Ahsan et al., 2009).

According to a survey across Bangladesh, As and other trace elements in paddy soil systems were elevated in areas where shallow tube wells had operated for a long time (Meharg and Rahman, 2003). However, microlevel studies in different villages of Chakdaha, Nadia, West Bengal, India, were conducted and confirmed the transference of As from irrigation water to the paddy soil-rice system (Bhattacharya et al., 2009). The results showed that the As concentration in the irrigation water ranged from 0.1 to 0.8 mg/L, which significantly exceeded the WHO acceptable limit (0.01 mg/L) for drinking and irrigation water; in the paddy soils, the reported As concentration was below the WHO acceptable limit (10 mg/kg). In rice plants, the maximum accumulation in roots ranged from 7.2 to 18.6 mg/kg, and the lowest accumulation, in the rice grains, ranged from 0.25 to 0.7 mg/kg (Arunakumara et al., 2013; Bhattacharya et al., 2009). A recent study conducted by Yadav et al. (2017) in the Fatehabad district, Haryana, India, suggested that trace metals (Cu, Co, & Zn) were abundant in rice grains as essential trace nutrients, and human health risk indices revealed that residents in most of the studied locations were safe. In Shakargarh, Pakistan, a study conducted by Tariq and Rashid (2012) reported Pb accumulation in rice grains, which may cause several health problems for the residents of the studied location.

Rahman et al. (2014) performed a comparative study of Australia-grown rice with market-purchased rice from India, Bangladesh, Pakistan, Canada, Italy, Thailand, and Vietnam from Asian shops in Sydney. Based on the results for Cu and Ni concentrations in Vietnamese rice and those for Cu, Ni, Cd, Pb, & Cr in Thai rice for sale in Australia, the metal concentrations exceeded those in the Australia-grown rice, whereas the trace metal concentrations in the Pakistani rice on sale in Australia were significantly lower than those in the Australia-grown rice.

6. Trace element accumulation influences in paddy soil-rice systems

Trace element accumulation in paddy systems is determined by several influences, such as the bioavailability of the metals, the physicochemical properties of the paddy soil, and the plant species of interest

(Kumarathilaka et al., 2018). According to Zhao et al. (2010), trace element enrichment in paddy plants is significantly correlated with the soil physicochemical properties, such as the soil pH and the soil organic matter (OM). However, the Cu accumulation in paddy plants showed a weak correlation with soil physicochemical properties, particularly the soil pH, and there was no correlation between the accumulation of Cu and OM. This suggests that other unknown factors could influence Cu accumulation in plants. A study conducted in the southern part of the Yangtze delta studied Cu contamination in the paddy soil-rice system, where Cu-rich contaminated wastewater is primarily used for irrigation purposes (Cao and Hu, 2000). Their results found a strong correlation between trace element concentrations in the soil and in rice grains, and the total Cu concentrations in the paddy soil, brown rice, rice hulls, and shoots were 101.2, 15.5, 133.2, and 101.4 mg/kg, respectively. Thus, strong correlations between the Cu concentration in rice shoots and rice hulls and the Cu concentration in the soil were proposed, with values of $r = 0.92$ & 0.83 , respectively (Cao and Hu, 2000).

According to Tariq and Rashid (2012), the trace element concentrations in paddy soil are significantly correlated with plant uptake but insignificantly correlated with the physicochemical properties of paddy soils and the plant species. Therefore, through plant rhizosphere processes, trace element uptake is generally governed by the physicochemical properties of paddy soil. According to them, the paddy soil pH was strongly negatively but insignificantly correlated with the paddy soil levels of Co, Pb, and Cr, with values of $r = -0.49$, -0.41 & -0.48 , respectively, whereas sulfate (SO_4^{2-}) was found in the paddy soils. However, the SO_4^{2-} content of the soil was strongly positively and significantly correlated with the Co, Pb, and Cr concentrations. Furthermore, significant positive correlations between pairs of trace elements in paddy soils, such as Pb and Cr, Cd and Pb, Pb and Co, and Cr and Co, and strong negative correlations between Ni and Cd as well as Pb and Ni were also reported. Nevertheless, within the plant parts or the grain, the relationships between trace elements were reported to be dissimilar from those in the paddy soil.

In Mae Sot, Tak, Thailand, no significant correlation was observed between the Cd concentration in paddy soil and that in rice grains, with $r = 0.117$ (Simmons et al., 2008). Additionally, in e-waste recycling areas in Southeast China, no strong significant correlation was observed between trace element concentrations in paddy soils and rice grains; furthermore, the investigators claimed that uptake may significantly contribute to Cd, Zn, & Pb levels, while Cu and Cr levels were primarily influenced by the roots (Fu et al., 2008). In Table 3, trace element translocation in the paddy soil-rice system (from the paddy soil to the paddy plant roots, shoots, and rice grains) is expressed via translocation factors (TFs) previously described by (Arunakumara et al., 2013; Gupta et al., 2008). The TF from the paddy soil to the grain ($TF_{\text{grain/soil}}$) was < 1 for all trace elements listed; nevertheless, a particularly high TF from the paddy soil to the root ($TF_{\text{root/soil}}$) is often shown (Arunakumara et al., 2013).

Trace elements such as Cd, Zn, & Pb and their particle size and mobility might be the significant reason for the maximum foliar and plant root uptake (Shahid et al., 2017b). It has been stated that the particle sizes of vehicle traffic-related elements such as Cd and Pb are usually fine, Cu and Zn are usually medium, and Cr is found in coarse particles; Zn is more mobile than Cu (Arunakumara et al., 2013). It has also been reported that Cd, Zn, and Pb particles emitted from vehicles have a higher leaf penetration capacity than those of Cu and Cr (Feng et al., 2011). Upon foliar Cd and Pb absorption, these metals might move to existing paddy plant growth areas, such as rice grains (Arunakumara et al., 2013). Thus, studies have suggested that trace element levels in rice grains not only depend on the physicochemical properties and available metal concentrations in the paddy soil system but also on vehicle emissions, the metal particle size, metal mobility, and metal penetration capability, which are significant factors if rice is grown adjacent to highways (Arunakumara et al., 2013).

7. Trace element-contaminated soil management/remediation methods

Once trace elements are released into the environment, they persist for an extended time period that varies depending on the amount and type of the trace element and the soil (Wuana and Okieimen, 2011). Several soil clean-up processes, such as physical, chemical, phytoremediation, and microbial remediation methods, are widely used in the remediation of trace elements. These methods can be performed *in situ* (on-site) or *ex situ* (off-site) (Fig. 4). These bioremediation methods are mostly used in combination for the cost-efficient and effective clean-up of trace element polluted sites (Liu et al., 2018).

7.1. Physical methods

Physical soil clean-up methods can be further classified into soil replacement, soil isolation, soil vitrification, and electrokinetic methods (Liu et al., 2018; Xia et al., 2019). In the soil replacement method, the polluted soil is replaced or partially replaced with non-contaminated soil (Liu et al., 2018). Before 1984, excavation, *ex situ*/off-site disposal, and soil replacement were the most common methods used for cleaning up contaminated sites (Khalid et al., 2017). Soil replacement dilutes the concentration of trace elements and, in turn, increases soil functionality (Wuana and Okieimen, 2011). In this method, the replaced soil is usually treated to remove trace elements or, in some cases, might be dumped at another site. The soil replacement method is generally carried out in one of two ways: digging the soil or importing new (uncontaminated) soil (Yao et al., 2012). In the digging method, the contaminated site is deeply excavated, and the trace elements are spread into the deeper places in the site. The trace elements are thereby diluted. The importing method refers to the addition of new, clean soil into the contaminated soil. The new clean soil might be used to cover the site surface or might be mixed in to decrease the levels of trace elements (Nejad et al., 2018; Kendall et al., 2017).

The soil isolation method refers to separating trace metal-polluted soil from uncontaminated soil; nonetheless, for complete remediation,

another auxiliary engineering method is needed (Marques et al., 2009). The soil isolation method is usually designed to stop the off-site movement of trace elements by limiting them to within a specified zone (Wuana and Okieimen, 2011; Marques et al., 2011). In soil, the isolation method is generally used to avoid further contamination of groundwater through trace elements after other remediation methods are not economically or physically possible (Liu et al., 2018). However, in some cases, the polluted zones can be isolated provisionally to prevent transportation during site evaluation and remediation (Khalid et al., 2017). In this method, the subsurface barriers/walls are typically applied to restrict polluted water and the soil by limiting the flow of both surface and groundwater in a contaminated site (Caliman et al., 2011).

In soil vitrification, the movement of trace elements inside the soil system is decreased through high-temperature action in highly contaminated sites, which leads to the formation of vitreous substances (Gong et al., 2018). In the soil vitrification method, some trace elements such as Hg can be volatilized under high-temperature conditions; therefore, they must be collected for additional disposal and/or further treatment (Rodríguez et al., 2012). The soil vitrification method is not considered a conventional trace element remediation method because the soil vitrification technique is relatively effortless to apply compared to other physical clean-up methods. Soil vitrification might be functional for most soils, which are polluted with both inorganic and organic pollutants (Khalid et al., 2017). Throughout the *in situ* soil vitrification method, an electrical current is passed through the soil by vertically inserting an array of electrodes into the polluted zone (Chen et al., 2016).

In contrast, the electrokinetic physical soil clean-up method is cost-effective and extensively used for the removal of trace elements from contaminated soils (Habibul et al., 2016). The soil electrokinetic process generally operates based on the electric field gradient principle with an appropriate power intensity and is typically applied to two sides of an electrolytic tank comprising saturated polluted soils (Vocciante et al., 2016). Trace elements that are present in the soil system can be separated through electrophoresis, electromigration, or




Fig. 4. Comparison of soil management/remediation methods for trace elements-contaminated soils.

electroosmosis, which decreases the contamination level of the contaminated site (Gong et al., 2018).

7.2. Chemical methods

Chemical-based soil clean-up methods can be further characterized as soil immobilization, soil encapsulation, and soil washing (Liu et al., 2018). The soil immobilization method refers to a reduction in trace element mobility, bioavailability, and bioaccessibility in the soil by adding immobilization agents to the polluted soil sites. In soil, trace elements are immobilized by several reactions, such as adsorption, complexation, and precipitation (Bolan et al., 2014). These immobilization reactions cause the redistribution of trace elements from the soil solution into solid particles, therefore reducing their movement and bioavailability in polluted soil. Trace element immobilization in the soil is mostly carried out by using inorganic and organic amendments to soils (Lwin et al., 2018). Common amendments to soils include microbes, phosphates, zeolites, cement, clays, minerals, and organic amendments (Khalid et al., 2017). Recently, several studies suggested that low-cost industrial residues, including industrial eggshells, red mud, and termitaries, could also be used for the immobilization of trace elements in contaminated soil (Khalid et al., 2017; Soares et al., 2015; Smičiklas et al., 2014). Currently, biomaterials are extensively used to immobilize trace elements in soils due to their availability and very low cost (Zhao et al., 2015a). Of the biomaterials, the extensive application of biochar has received important consideration for immobilizing trace elements in the soil (Cao et al., 2011).

The soil encapsulation method differs from the other methods and is mostly used as a process to filter contaminants from soils; the encapsulation stops them from spreading further (Caliman et al., 2011). Soil encapsulation remediation also involves the mixing of polluted soils with different products, such as concrete, lime, and asphalt. (Wuana and Okieimen, 2011; Khalid et al., 2017). In the encapsulation method, the toxic element-contaminated soil becomes immobile and therefore stops the spreading of pollution to surrounding resources (France, 2015). Several binding materials are used in the formation of the solid blocks; cement is the ideal material due to its high availability, adaptability, and cost-effectiveness (Wuana and Okieimen, 2011; Khalid et al., 2017).

The soil washing method refers to the removal of trace elements from contaminated soils by using different reagents and extractants that can leach trace elements from the soil. It is done on physically excavated soils *ex situ* or *in situ* (Wuana and Okieimen, 2011; Dermont et al., 2008). Recently, the application of appropriate extractants for the leaching of trace metals from polluted soils has been confirmed as an excellent alternative to some of the old conventional methods for the removal of toxic elements (Wuana and Okieimen, 2011; Leštan et al., 2008). During the soil washing method, the polluted soil is excavated and mixed with an appropriate extractant solution based on the type of trace elements and soil properties (Begum et al., 2016).

7.3. Phytoremediation

Phytoremediation is a practical remediation method and can rebuild the natural conditions of the soil to remediate harmful or adverse ecological effects (Apori et al., 2018). Phytoremediation methods primarily make use of plants to remove contaminants from soils (Chibuike and Obiora, 2014). The phytoremediation method is comparatively efficient, simple, non-invasive, cost-effective, and eco-friendly and can provide an enduring solution (Sivakumar et al., 2014). Phytoremediation is also known as green remediation, botanoremediation, agrorremediation, and/or vegetative remediation and includes methods that generally make use of plants to remediate and revegetate contaminated sites (Wuana and Okieimen, 2011). The term phytoremediation includes several methods and applications that differ significantly in process or mechanisms with plants that might immobilize, remove/

clean-up, or degrade toxic trace elements (Yadav et al., 2018). Based on diverse plant uptake patterns, phytoremediation can include phytostabilization, photoevaporation, and phytoextraction (Awa and Hadibarata, 2020). During phytovolatilization, toxic trace elements are taken up from the soil and transformed into less-toxic vapours and then released into the atmosphere via the plant transpiration process (Singh et al., 2017).

During the phytovolatilization process, elements are first combined into volatile organic complexes and then released to the ambient environment as biomolecules (Kumar and Gunasundari, 2018). Some trace elements, including As, Se, and Hg, may occur as gases in the environment. Several plant species, such as *Brassica juncea*, *Arabidopsis thaliana*, and *Chara canescens*, have the capability to take up these toxic elements from soils and change them into their gaseous forms inside the plant tissues and then release them into the ambient environment. Arsenic and its species As(V) and As(III) were effectively volatilized from the leaves of *Pteris vittata* plants. *Arabidopsis thaliana* and *Brassica juncea* can be cultivated under elevated Se levels and are proficient in volatilizing Se (Khalid et al., 2017).

Phytostabilization is well defined as the use of plants to decrease the availability and movement of toxic elements in soils because of their stabilization by plants (Mahar et al., 2016). Phytostabilization may not reduce the levels of toxic elements that exist in polluted soil but prevents their off-site mobilization (Sarwar et al., 2017). The primary purpose of phytostabilization is to limit trace elements in the vadose zone of plants by accumulation through roots and/or precipitation inside the rhizosphere zone (Bolan et al., 2011; Pinto et al., 2015). Thus, of the different phytoremediation methods, phytostabilization does not remediate or remove toxic elements from soils but decreases the mobilization of toxic elements to adjacent media/sites (Sarwar et al., 2017). In addition, phytostabilization may also be applied at contaminated sites where practical/controlling limits interfere with the selection and application of the most suitable clean-up methods (Tack and Meers, 2010). Several plant species, for instance, *Agrostis* spp., and *Festuca* spp. are widely applied for the phytostabilization of Cu-, Pb-, and Zn-contaminated soils in Europe (Galende et al., 2014).

Phytoextraction is an efficient method for the removal of toxic elements from contaminated soils through plant uptake mechanisms (Laghlimi et al., 2015). This solar-driven method is generally based on the ability of plant roots to take up and translocate levels of toxic elements from the soil to the above-ground plant parts (Ali et al., 2013). Therefore, phytoextraction reduces the contaminated soil mass (Wan et al., 2016) and efficiently translocates toxic elements from the soil to aerial plant parts (Bhargava et al., 2012). Compared with contaminated soil, plant biomass is generally easy to recycle, transfer, and treat/oxidize, and this method ensures the a permanent removal of toxic elements from soils (Sarwar et al., 2017). Nevertheless, the phytoextraction method is more convenient for those sites that are contaminated by low-to-moderate levels of toxic elements because several plant species may not be able to survive in highly contaminated sites (Sarwar et al., 2017; Hashim et al., 2011). For instance, the phytoextraction method is very efficient for Pb removal with *Brassica juncea* in soils whose Pb concentration is less than 1500 mg/kg (Khalid et al., 2017).

Typically, plant species that are efficiently used for phytoextraction must have 1) high toxic element accumulation ability in their aerial parts, 2) high tolerance to levels of toxic elements, 3) high capability to grow quickly with maximum biomass, and 4) very profuse root systems; so far, no standard has been suggested for hyperaccumulator plants (Khalid et al., 2017). However, hyperaccumulator plant species can accumulate 100- to 500-fold more toxic elements in their shoot tissues than non-accumulator plants without any effect on yield (Khalid et al., 2017; Chaney et al., 2007). Typically, most hyperaccumulator plant species can accumulate or tolerate only a single specific toxic element (Clemens, 2006), while fewer plant species have the ability to accumulate and tolerate more than one element; these plants are known as

multielement hyperaccumulator plant species (Padmavathiamma and Li, 2007). Examples include *Thlaspi caerulescens*, which hyperaccumulates Pb, Zn, Cd, and Ni (Ali et al., 2013); *Thlaspi goesingense* and *Thlaspi ochroleucum* can hyperaccumulate Ni and Zn; and *Thlaspi rotundifolium* efficiently hyperaccumulates Pb, Zn, and Ni (Khalid et al., 2017).

Phytoextraction efficiency is increased through genetic modification of plants (da Conceição Gomes et al., 2016). It has been suggested that several genes play a crucial role in toxic element uptake, translocation, and sequestration in the vacuole in plant tissues (Thakur et al., 2016). Primarily, the application of a gene in specific plant species typically depends on the objective of plant mutation (Yang et al., 2005). In phytoextraction processes, enhanced uptake with an increased translocation of elements towards the plant's aerial parts (Sarwar et al., 2017) can be accomplished through genetically modified genes that are mainly responsible for enhancing element adsorption in plant root cells, decreasing plant root element sequestration, increasing element xylem loading, and increasing translocation to plant shoots for storage in epidermal and mesophyll cells (Khalid et al., 2017). To date, numerous key processes have been reported at the molecular level, signifying the possibility of engineering toxic element hyperaccumulation in plant systems (Kumar et al., 2017). The multifold enhancement in absorption, xylem loading, and translocation to aerial shoots increases the tolerance reported in genetically modified plants compared to that in wild types (Khalid et al., 2017) for some toxic elements, such as As, Zn, Pb, and Cd, in *Arabidopsis thaliana*, *Lactuca sativa*, *Brassica oleracea*, *Nicotiana glauca*, and *N. tabacum*, respectively (Guo et al., 2012; Barrameda-Medina et al., 2014; Khoudi et al., 2013; Gisbert et al., 2003).

7.4. Microbial remediation

Microbial remediation primarily refers to the application of microorganisms to promote the precipitation, adsorption, and oxidation-reduction of trace elements in the soil. Under toxic metal stress conditions, soil microbial associations aid in increasing plant growth (Ayangbenro and Babalola, 2017). Several studies have reported that plant growth-promoting and toxic element resistance and soil microorganisms might also protect plants from element toxicity and/or even increase element uptake in hyperaccumulator plants (Pinto et al., 2018). Microorganism-based remediation methods for toxic elements involve numerous mechanisms, including bioleaching, biomineralization, biosorption, enzyme-catalysed transformation, intracellular accumulation, and oxidation-reduction reactions (Khalid et al., 2017). Toxic element-resistant rhizobacteria are able to stimulate plant growth and element accumulation in plants by the production of different chemicals, such as indole acetic acid (IAA) and monocyclopropane-1-carboxylate (ACC) (Glick, 2014). Furthermore, element-resistant rhizobacteria increase the element tolerance of host plants by inducing thiol complexes, metallothionein, and/or superoxide dismutase (Khalid et al., 2017; Cuyppers et al., 2013). Application of *Burkholderia* sp. (Z-90) increased toxic element clean-up efficiency in the soil by 24 %, 31 %, 32 %, 37 %, 44 %, and 52 % for Cu, Pb, As, Cd, Zn, and Mn, respectively (Yang et al., 2016). *Bacillus* sp. (E1S2) significantly enhanced the uptake of Zn (18 %) and Cd (43 %) by *Sedum plumbizincicola* (Ma et al., 2015). In addition, microbial associations may also cause numerous positive effects for plant growth, including osmotic adjustment, stomatal regulation, and increased mineral uptake (Ma et al., 2011). Soil bacterial associations may also increase the mobility and availability of the elements in soils in numerous ways, including decreasing the soil pH, facilitating plant-growth (Tak et al., 2013), and chelating complexes (organic acids, biosurfactants, and siderophores) through changing the soil oxidation-reduction conditions (Khalid et al., 2017). Nevertheless, the influence of soil microorganisms on element solubility and plant uptake may not be consistent (de la Fuente et al., 2011). Soil microorganisms might increase or decrease element uptake by plants (Sessitsch et al., 2013). Indeed, specific microbial methods can increase

element solubility and bioaccumulation; however, numerous other methods may cause element immobilization, therefore reducing their phytoextraction (Khalid et al., 2017; Pinto et al., 2018).

Additionally, several other important factors that influence the selection and applicability of the existing soil remediation processes include the total cost involved, time required, efficiency under elevated toxic element contamination, acceptance and availability, long-term effectiveness, and applicability to multi-element contaminated sites; these factors are discussed in detail in Table S3 (Khalid et al., 2017).

8. Conclusion and future research perspectives

In this review, we summarize the basic chemical behaviours, transport, endpoints, and significant natural and anthropogenic sources of trace elements, including As, Se, Cd, Pb, Zn, Mn, Ni, and Cr, in the soil system as well as their uptake, translocation, and accumulation mechanisms in the paddy soil-rice system. Human health effects are significantly associated with these toxic and potentially toxic metals not only due to their occurrence in the environment but also in their uptake, translocation, and accumulation via different transporters, such as As (silicate (Si), phosphate (Pi)), Se (sulfate (ST)), Cr (Pi or ST), Zn (AtZIP1 or NRAMP), Mn (NRAMP5 or AtZIP1), Ni (NRAMP), and Cd (NRAMP5, OsIRT1, or osHMA2), in rice systems.

Moreover, the contamination status of paddy soil-rice systems in major rice-growing countries, including India, Bangladesh, China, Pakistan, Malaysia, Thailand, Japan, and Australia, was presented. Consequently, it is worth comparing the existing research conclusions on the elevated concentrations of trace elements in rice grain in different countries: Se, As, and Zn (0.02–9.7, 3.1–14, & 2.3–484.9), Cd, and Pb (9.4–178, & 0.06–1529), Cr (41–217), and Ni (61–356) in China, India, Bangladesh, and Australia, respectively.

This review also provides an inventory of the various research studies on trace metal concentrations from multiple identified and unidentified contamination sources, mainly in paddy soil-rice systems in rice-growing countries. In the future, this study will be helpful for paddy rice farmers and decisionmakers at regional, national, and international levels to support various administrative policies. The biogeochemical conditions associated with paddy farming, such as the frequent alternation between wet and dehydrated environments and the oxygenated-non oxygenated paddy soils, make paddy rice systems favourable for trace metal accumulation.

The various influences associated with the accumulation of these metals in the paddy soil-rice system could be 1) the paddy soil-rice plant characteristics, 2) the various soil physico-chemical indicators associated with uptake, translocation, and accumulation, and 3) the post-crop management practices. These mechanisms are primarily reliant on the cultivar (product class), the environmental conditions, and the administrative policies. Accordingly, multidisciplinary elementary and practical research efforts are essential to mitigate trace metal accumulation in paddy soil-rice systems and prevent toxicity to humans from rice.

The contents of the current review highlight the need for a wide-ranging, extensive, periodic monitoring of the contamination status of paddy soil-rice system regions. Broad awareness of paddy soil contamination and the associated health risks might lead different countries to establish strict legal guidelines and rules for paddy soil contamination and to adopt consistent remediation measures to control trace element sources and translocation in paddy soil-rice systems.

Therefore, the present study will be helpful for comparing the various trace element mitigation measures and for understanding the status of trace element contamination, categorization, and translocation mechanisms in paddy soil-rice systems.

Declaration of Competing Interest

The authors declared no conflict of interest.

Acknowledgments

This work is supported by National Key Research and Development Program of China (Nos. 2019 YFC1803600, and 2017YFD0800302), the National Science Foundation of China-Project of Karst Science Research Center (U1612442), and Science and Technology program of Guizhou Province [Grant No. (2019)5618/2019(2963)/(2019)4428], UK Royal Academy of Engineering Frontier Follow-up Grant (FF1920136). The authors would also like to thank all anonymous reviewers for their constructive comments that significantly improved the quality of the manuscript.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.jhazmat.2020.122720>.

References

- Abbas, S.H., Ismail, I.M., Mostafa, T.M., Sulaymon, A.H., 2014. Biosorption of heavy metals: a review. *J. Chem. Sci. Technol.* 3, 74–102.
- Abbas, G., Murtaza, B., Bibi, I., Shahid, M., Niazi, N., Khan, M., Amjad, M., Hussain, M., 2018. Arsenic uptake, toxicity, detoxification, and speciation in plants: physiological, biochemical, and molecular aspects. *Int. J. Environ. Res. Public Health* 15, 59.
- Abraham, J., Dowling, K., Florentine, S., 2018. Assessment of potentially toxic metal contamination in the soils of a legacy mine site in Central Victoria, Australia. *Chemosphere* 192, 122–132.
- Abtahi, M., Fakhri, Y., Oliveri Conti, G., Keramati, H., Zandsalimi, Y., Bahmani, Z., Hosseini Pouya, R., Sarkhosh, M., Moradi, B., Amanidaz, N., 2017. Heavy metals (As, Cr, Pb, Cd and Ni) concentrations in rice (*Oryza sativa*) from Iran and associated risk assessment: a systematic review. *Toxin Rev.* 36, 331–341.
- Adriano, D.C., 2013. *Trace Elements in the Terrestrial Environment*. Springer Science & Business Media.
- Ahmadi, H., 2017. Functional Analysis of the Metal Hyperaccumulation and Hypertolerance Candidate Genes NAS4, ZIP6, CAX1 and NRAMP3 in Arabidopsis halleri. Universität Bayreuth.
- Ahsan, D.A., DelValls, T.A., Blasco, J., 2009. Distribution of arsenic and trace metals in the floodplain agricultural soil of Bangladesh. *Bull. Environ. Contam. Toxicol.* 82, 11–15.
- Ali, H., Khan, E., Sajad, M.A., 2013. Phytoremediation of heavy metals—concepts and applications. *Chemosphere* 91, 869–881.
- Ali, W., Aslam, M.W., Feng, C., Junaid, M., Ali, K., Li, S., Chen, Z., Yu, Z., Rasool, A., Zhang, H., 2019a. Unraveling prevalence and public health risks of arsenic, uranium and co-occurring trace metals in groundwater along riverine ecosystem in Sindh and Punjab, Pakistan. *Environ. Geochem. Health* 41, 2223–2238.
- Ali, W., Mushtaq, N., Javed, T., Zhang, H., Ali, K., Rasool, A., Farooqi, A., 2019b. Vertical mixing with return irrigation water the cause of arsenic enrichment in groundwater of district Larkana Sindh, Pakistan. *Environ. Pollut.* 245, 77–88.
- Ali, W., Zhang, H., Junaid, M., Mao, K., Xu, N., Chang, C., Rasool, A., Wajhat Aslam, M., Ali, J., Yang, Z., 2020. Insights into the mechanisms of arsenic-selenium interactions and the associated toxicity in plants, animals, and humans: a critical review. *Crit. Rev. Environ. Sci. Technol.* 1–47.
- Alloway, B.J., 2013a. Sources of heavy metals and metalloids in soils. *Heavy Metals in Soils*. Springer, pp. 11–50.
- Alloway, B.J., 2013b. Heavy metals and metalloids as micronutrients for plants and animals. *Heavy Metals in Soils*. Springer, pp. 195–209.
- Antoniadis, V., Levizou, E., Shaheen, S.M., Ok, Y.S., Sebastian, A., Baum, C., Prasad, M.N., Wenzel, W.W., Rinklebe, J., 2017. Trace elements in the soil-plant interface: Phytoavailability, translocation, and phytoremediation—a review. *Earth. Rev.* 171, 621–645.
- Antoniadis, V., Shaheen, S.M., Levizou, E., Shahid, M., Niazi, N.K., Vithanage, M., Ok, Y.S., Bolan, N., Rinklebe, J., 2019a. A critical prospective analysis of the potential toxicity of trace element regulation limits in soils worldwide: are they protective concerning health risk assessment? A review. *Environ. Int.* 127, 819–847.
- Antoniadis, V., Golia, E.E., Liu, Y.-T., Wang, S.-L., Shaheen, S.M., Rinklebe, J., 2019b. Soil and maize contamination by trace elements and associated health risk assessment in the industrial area of Volos, Greece. *Environ. Int.* 124, 79–88.
- Apori, O., Hanyabui, E., Asiamah, Y., 2018. Remediation technology for copper contaminated soil: a review. *Asian Soil Res. J.* 1–7.
- ar Hashmi, M.Z., Varma, A., 2018. Environmental Pollution of Paddy Soils.
- Arao, T., Ishikawa, S., Murakami, M., Abe, K., Maejima, Y., Makino, T., 2010. Heavy metal contamination of agricultural soil and countermeasures in Japan. *Paddy Water Environ.* 8, 247–257.
- Arroyo, M.d.M.D., Hornedo, R.M.D.I., Peralta, F.A., Almestre, C.R., Sánchez, J.V.M., 2014. Heavy metals concentration in soil, plant, earthworm and leachate from poultry manure applied to agricultural land. *Rev. Int. Contam. Ambient.* 30, 43–50.
- Arunakumara, K., Walpola, B.C., Yoon, M.-H., 2013. Current status of heavy metal contamination in Asia's rice lands. *Rev. Environ. Sci. Biotechnol.* 12, 355–377.
- Asgari, F., Haghazali, M., Heydarian, H., 2009. Non-Communicable Diseases Risk Factors Surveillance in Iran.
- Awa, S.H., Hadibarata, T., 2020. Removal of heavy metals in contaminated soil by phytoremediation mechanism: a review. *Water Air Soil Pollut.* 231, 47.
- Awan, K.Y., Rashid, A., 2012. Overview of Pakistan's electricity crisis, generation-mix and renewable energy scenarios. *Int. J. Eng. Technol.* 1, 321–334.
- Ayangbenro, A.S., Babalola, O.O., 2017. A new strategy for heavy metal polluted environments: a review of microbial biosorbents. *Int. J. Environ. Res. Public Health* 14, 94.
- Babayemi, J.O., Ogundiran, M.B., Osibanjo, O., 2016. Overview of environmental hazards and health effects of pollution in developing countries: a case study of Nigeria. *Environ. Qual. Manag.* 26, 51–71.
- Bai, J., Cui, B., Wang, Q., Gao, H., Ding, Q., 2009. Assessment of heavy metal contamination of roadside soils in Southwest China. *Stoch. Environ. Res. Risk Assess.* 23, 341–347.
- Barrameda-Medina, Y., Montesinos-Pereira, D., Romero, L., Ruiz, J.M., Blasco, B., 2014. Comparative study of the toxic effect of Zn in *Lactuca sativa* and *Brassica oleracea* plants: I. Growth, distribution, and accumulation of Zn, and metabolism of carboxylates. *Environ. Exp. Bot.* 107, 98–104.
- Bastías, J.M., Beldarrain, T., 2016. Arsenic translocation in rice cultivation and its implication for human health. *Chil. J. Agric. Res.* 76, 114–122.
- Baysal, A., Ozbek, N., Akman, S., 2013. Determination of trace metals in waste water and their removal processes. In: *Einschlag, F.S.G., Carlos, L. (Eds.), Waste-Water Treatment Technologies and Recent Analytical Developments*, pp. 145–171.
- Begum, Z.A., Rahman, I.M., Sawai, H., Hasegawa, H., 2016. Chemical-induced washing remediation of metal-contaminated soils. *Environmental Remediation Technologies for Metal-Contaminated Soils*. Springer, pp. 197–218.
- Bhargava, A., Carmona, F.F., Bhargava, M., Srivastava, S., 2012. Approaches for enhanced phytoextraction of heavy metals. *J. Environ. Manage.* 105, 103–120.
- Bhattacharya, P., Mukherjee, A.B., Bundschuh, J., Zevenhoven, R., Loeppert, R.H., 2007. *Arsenic in Soil and Groundwater Environment: Biogeochemical Interactions, Health Effects and Remediation*. Elsevier.
- Bhattacharya, P., Samal, A., Majumdar, J., Santra, S., 2009. Transfer of arsenic from groundwater and paddy soil to rice plant (*Oryza sativa* L.): a micro level study in West Bengal, India. *World J. Agric. Sci.* 5, 425–431.
- Bini, C., 2013. Geology and geomorphology. *The Soils of Italy*. Springer, pp. 39–56.
- Biswas, S., Mojid, M., Wyseure, G., 2017. Physicochemical properties of soil under wheat cultivation by irrigation with municipal wastewater in Bangladesh. *Commun. Soil Sci. Plant Anal.* 48, 1–10.
- Bitterli, C., Bañuelos, G., Schulin, R., 2010. Use of transfer factors to characterize uptake of selenium by plants. *J. Geochem. Explor.* 107, 206–216.
- Bolan, N.S., Park, J.H., Robinson, B., Naidu, R., Huh, K.Y., 2011. Phytostabilization: a green approach to contaminant containment. *Advances in Agronomy*. Elsevier, pp. 145–204.
- Bolan, N., Kunhikrishnan, A., Thangarajan, R., Kumpiene, J., Park, J., Makino, T., Kirkham, M.B., Scheckel, K., 2014. Remediation of heavy metal (loid)s contaminated soils—to mobilize or to immobilize? *J. Hazard. Mater.* 266, 141–166.
- Borthakur, A., Govind, M., 2017. Emerging trends in consumers' E-waste disposal behaviour and awareness: a worldwide overview with special focus on India. *Resources. Conserv. Recycl.* 117, 102–113.
- Bretzel, F.C., Calderisi, M., 2011. Contribution of a municipal solid waste incinerator to the trace metals in the surrounding soil. *Environ. Monit. Assess.* 182, 523–533.
- Brown, K.G., Ross, G.L., 2002. Arsenic, drinking water, and health: a position paper of the American Council on Science and Health. *Regul. Toxicol. Pharmacol.* 36, 162–174.
- Cai, L., Huang, L., Zhou, Y., Xu, Z., Peng, X., Yao, L.a., Zhou, Y., Peng, P.a., 2010. Heavy metal concentrations of agricultural soils and vegetables from Dongguan, Guangdong. *J. Geogr. Sci.* 20, 121–134.
- Caliman, F.A., Robu, B.M., Smaranda, C., Pavel, V.L., Gavrilescu, M., 2011. Soil and groundwater cleanup: benefits and limits of emerging technologies. *Clean Technol. Environ. Policy* 13, 241–268.
- Campbell, K.M., Nordstrom, D.K., 2014. Arsenic speciation and sorption in natural environments. *Rev. Mineral. Geochem.* 79, 185–216.
- Candeias, C., Melo, R., Ávila, P.F., da Silva, E.F., Salgueiro, A.R., Teixeira, J.P., 2014. Heavy metal pollution in mine-soil-plant system in S. Francisco de Assis-Panasqueira mine (Portugal). *Appl. Geochem.* 44, 12–26.
- Cao, Z., Hu, Z., 2000. Copper contamination in paddy soils irrigated with wastewater. *Chemosphere* 41, 3–6.
- Cao, Z., Wang, X., Yao, D., Zhang, X., Wong, M., 2001. Selenium geochemistry of paddy soils in Yangtze River Delta. *Environ. Int.* 26, 335–339.
- Cao, H., Chen, J., Zhang, J., Zhang, H., Qiao, L., Men, Y., 2010. Heavy metals in rice and garden vegetables and their potential health risks to inhabitants in the vicinity of an industrial zone in Jiangsu, China. *J. Environ. Sci.* 22, 1792–1799.
- Cao, X., Ma, L., Liang, Y., Gao, B., Harris, W., 2011. Simultaneous immobilization of lead and atrazine in contaminated soils using dairy-manure biochar. *Environ. Sci. Technol.* 45, 4884–4889.
- Chandrakar, V., Pandey, N., Keshavkant, S., 2018. Plant responses to arsenic toxicity: morphology and physiology. *Mechanisms of Arsenic Toxicity and Tolerance in Plants*. Springer, pp. 27–48.
- Chaney, R.L., Angle, J.S., Broadhurst, C.L., Peters, C.A., Tappero, R.V., Sparks, D.L., 2007. Improved understanding of hyperaccumulation yields commercial phytoextraction and phytomining technologies. *J. Environ. Qual.* 36, 1429–1443.
- Chang, C., Yin, R., Wang, X., Shao, S., Chen, C., Zhang, H., 2019. Selenium translocation

- in the soil-rice system in the Enshi seleniferous area, Central China. *Sci. Total Environ.* 669, 83–90.
- Chauhan, R., Awasthi, S., Srivastava, S., Dwivedi, S., Pilon-Smits, E.A., Dhankher, O.P., Tripathi, R.D., 2019. Understanding selenium metabolism in plants and its role as a beneficial element. *Crit. Rev. Environ. Sci. Technol.* 1–22.
- Chen, T., Liu, X., Zhu, M., Zhao, K., Wu, J., Xu, J., Huang, P., 2008. Identification of trace element sources and associated risk assessment in vegetable soils of the urban–rural transitional area of Hangzhou, China. *Environ. Pollut.* 151, 67–78.
- Chen, Z., Zhao, Y., Li, Q., Qiao, J., Tian, Q., Liu, X., 2009a. Heavy metal contents and chemical speciations in sewage-irrigated soils from the eastern suburb of Beijing, China. *J. Food Agric. Environ.* 7 (3).
- Chen, S., Sun, L., Chao, L., Zhou, Q., Sun, T., 2009b. Estimation of lead bioavailability in smelter-contaminated soils by single and sequential extraction procedure. *Bull. Environ. Contam. Toxicol.* 82, 43–47.
- Chen, H., Yan, S., Ye, Z., Meng, H., Zhu, Y., 2012. Utilization of urban sewage sludge: Chinese perspectives. *Environ. Sci. Pollut. Res. - Int.* 19, 1454–1463.
- Chen, J.P., Wang, L.K., Wang, M.-H.S., Hung, Y.-T., Shammass, N.K., 2016. Remediation of Heavy Metals in the Environment. *Crc Press*.
- Chen, Y., Han, Y.-H., Cao, Y., Zhu, Y.-G., Rathinasabapathi, B., Ma, L.Q., 2017. Arsenic transport in rice and biological solutions to reduce arsenic risk from rice. *Front. Plant Sci.* 8, 268.
- Cheng, X., Drozdova, J., Danek, T., Huang, Q., Qi, W., Yang, S., Zou, L., Xiang, Y., Zhao, X., 2018. Pollution assessment of trace elements in agricultural soils around copper mining area. *Sustainability* 10, 4533.
- Chiban, M., Zerbet, M., Carja, G., Sinan, F., 2012. Application of low-cost adsorbents for arsenic removal: a review. *J. Environ. Chem. Ecotoxicol.* 4, 91–102.
- Chibuike, G.U., Obiora, S.C., 2014. Heavy metal polluted soils: effect on plants and bioremediation methods. *Appl. Environ. Soil Sci.* 2014.
- Chinoim, N., Sinbuathong, N., 2010. Heavy metal contamination of soils from organic paddy fields in Thailand. 19th World Congress of Soil Science, Soil Solutions for a Changing World held on 1–6.
- Chu, T.T.H., 2011. Survey on Heavy Metals Contaminated Soils in Thai Nguyen and Hung Yen Provinces in Northern Vietnam.
- Cimen, O., Gok, G., Celebi, H., Gok, O., 2014. The possible environmental impacts of the biosolids in the worldwide. *J. Selcuk Univ. Nat. Appl. Sci.* 4, 1144–1154.
- Clarke, R., 2013. *Water: the International Crisis*. Routledge.
- Clemens, S., 2006. Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants. *Biochimie* 88, 1707–1719.
- Clemens, S., Ma, J.F., 2016. Toxic heavy metal and metalloid accumulation in crop plants and foods. *Annu. Rev. Plant Biol.* 67, 489–512.
- Cuyppers, A., Remans, T., Weyens, N., Colpaert, J., Vassilev, A., Vangronsveld, J., 2013. Soil-plant relationships of heavy metals and metalloids. *Heavy Metals in Soils*. Springer, pp. 161–193.
- da Conceição Gomes, M.A., Hauser-Davis, R.A., de Souza, A.N., Vitória, A.P., 2016. Metal phytoremediation: general strategies, genetically modified plants and applications in metal nanoparticle contamination. *Ecotoxicol. Environ. Saf.* 134, 133–147.
- Dang, T., Nguyen, P., Nguyen, D., 2008. Soil pollution with heavy metals in the some areas in Vietnam. *Vietnam Soil Sci. J.* 29, 59–62.
- de la Fuente, C., Clemente, R., Martínez-Alcalá, I., Tortosa, G., Bernal, M.P., 2011. Impact of fresh and composted solid olive husk and their water-soluble fractions on soil heavy metal fractionation; microbial biomass and plant uptake. *J. Hazard. Mater.* 186, 1283–1289.
- Dermont, G., Bergeron, M., Mercier, G., Richer-Lafleche, M., 2008. Soil washing for metal removal: a review of physical/chemical technologies and field applications. *J. Hazard. Mater.* 152, 1–31.
- Deshpande, J.D., Joshi, M.M., Giri, P.A., 2013. Zinc: the trace element of major importance in human nutrition and health. *Int. J. Med. Sci. Public Health* 2, 1–6.
- Dhillon, K.S., Dhillon, S.K., 2018. Genesis of Seleniferous Soils and Associated Animal and Human Health Problems.
- Dotaniya, M., Meena, V., 2015. Rhizosphere effect on nutrient availability in soil and its uptake by plants: a review. *Proc. Natl. Acad. Sci. India Sect. B: Biol. Sci.* 85, 1–12.
- Dotaniya, M., Thakur, J., Meena, V., Jajoria, D., Rathor, G., 2014. Chromium pollution: a threat to environment—a review. *Agric. Rev.* 35.
- Du Laing, G., Rinklebe, J., Vandecasteele, B., Meers, E., Tack, F.M., 2009. Trace metal behaviour in estuarine and riverine floodplain soils and sediments: a review. *Sci. Total Environ.* 407, 3972–3985.
- Dung, T.T.T., Cappuyns, V., Swennen, R., Phung, N.K., 2013. From geochemical background determination to pollution assessment of heavy metals in sediments and soils. *Rev. Environ. Sci. Biotechnol.* 12, 335–353.
- Dutta, S., Mitra, M., Agarwal, P., Mahapatra, K., De, S., Sett, U., Roy, S., 2018. Oxidative and genotoxic damages in plants in response to heavy metal stress and maintenance of genome stability. *Plant Signal. Behav.* 13, e1460048.
- Echevarria, G., 2018. Genesis and behaviour of ultramafic soils and consequences for nickel biogeochemistry. *Agromining: Farming for Metals*. Springer, pp. 135–156.
- Ehsan, M., Barakat, M., Husein, D.Z., Ismail, S., 2014. Immobilization of Ni and Cd in soil by biochar derived from unfertilized dates. *Water Air Soil Poll.* 225, 2123.
- Elgallal, M., Fletcher, L., Evans, B., 2016. Assessment of potential risks associated with chemicals in wastewater used for irrigation in arid and semiarid zones: a review. *Agric. Water Manage.* 177, 419–431.
- Essington, M.E., 2015. *Soil and Water Chemistry: an Integrative Approach*. CRC press.
- Farkas, E., Buglyó, P., 2017. Lead (II) complexes of amino acids, peptides, and other related ligands of biological interest. *Lead: Its Effects on Environment and Health*, vol. 17. pp. 201.
- Fendorf, S., Michael, H.A., van Geen, A., 2010. Spatial and temporal variations of groundwater arsenic in South and Southeast Asia. *Science* 328, 1123–1127.
- Feng, J., Wang, Y., Zhao, J., Zhu, L., Bian, X., Zhang, W., 2011. Source attributions of heavy metals in rice plant along highway in Eastern China. *J. Environ. Sci.* 23, 1158–1164.
- Ferrante, M., 2013. Manganese (Mn), Health Effects of Metals and Related Substances in Drinking Water. 71.
- Finnegan, P., Chen, W., 2012. Arsenic toxicity: the effects on plant metabolism. *Front. Physiol.* 3, 182.
- Fordeyce, F.M., 2013. Selenium deficiency and toxicity in the environment. *Essentials of Medical Geology*. Springer, pp. 375–416.
- France, C.B., **Method of encapsulating waste**, Google Patents, 2015.
- Fu, J., Zhou, Q., Liu, J., Liu, W., Wang, T., Zhang, Q., Jiang, G., 2008. High levels of heavy metals in rice (*Oryzasativa L.*) from a typical E-waste recycling area in southeast China and its potential risk to human health. *Chemosphere* 71, 1269–1275.
- Fujimaki, S., Suzui, N., Ishioka, N.S., Kawachi, N., Ito, S., Chino, M., Nakamura, S.-i., 2010. Trace cadmium from culture to spikelet: noninvasive imaging and quantitative characterization of absorption, transport, and accumulation of cadmium in an intact rice plant. *Plant Physiol.* 152, 1796–1806.
- Funari, V., Bokhari, S.N.H., Vigliotti, L., Meisel, T., Braga, R., 2016. The rare earth elements in municipal solid waste incinerators ash and promising tools for their prospecting. *J. Hazard. Mater.* 301, 471–479.
- Gailer, J., 2007. Arsenic–selenium and mercury–selenium bonds in biology. *Coord. Chem. Rev.* 251, 234–254.
- Galende, M.A., Becerril, J.M., Barrutia, O., Artetxe, U., Garbisu, C., Hernández, A., 2014. Field assessment of the effectiveness of organic amendments for aided phytostabilization of a Pb–Zn contaminated mine soil. *J. Geochem. Explor.* 145, 181–189.
- Garba, S., Abubakar, M., 2018. Source and distribution of the heavy metals: Pb, Cd, Cu, Zn, Fe, Cr, and Mn in soils of bauchi metropolis, Nigeria. *Am. J. Eng. Res.* 7, 13–23.
- Geng, N., Wu, Y., Zhang, M., Tsang, D.C., Rinklebe, J., Xia, Y., Lu, D., Zhu, L., Palansoriya, K.N., Kim, K.-H., 2019. Bioaccumulation of potentially toxic elements by submerged plants and biofilms: a critical review. *Environ. Int.* 131, 105015.
- Gisbert, C., Ros, R., De Haro, A., Walker, D.J., Bernal, M.P., Serrano, R., Navarro-Aviñó, J., 2003. A plant genetically modified that accumulates Pb is especially promising for phytoremediation. *Biochem. Biophys. Res. Commun.* 303, 440–445.
- Głąb, T., Żabiński, A., Sadowska, U., Gondek, K., Kopeć, M., Mierzwa-Hersztek, M., Tabor, S., 2018. Effects of co-composted maize, sewage sludge, and biochar mixtures on hydrological and physical qualities of sandy soil. *Geoderma* 315, 27–35.
- Glick, B.R., 2014. Bacteria with ACC deaminase can promote plant growth and help to feed the world. *Microbiol. Res.* 169, 30–39.
- Gong, Y., Zhao, D., Wang, Q., 2018. An overview of field-scale studies on remediation of soil contaminated with heavy metals and metalloids: technical progress over the last decade. *Water Res.* 147, 440–460.
- González-Corrochano, B., Esbrí, J., Alonso-Azcárate, J., Martínez-Coronado, A., Jurado, V., Higuera, P., 2014. Environmental geochemistry of a highly polluted area: the La Unión Pb–Zn mine (Castilla-La Mancha region, Spain). *J. Geochem. Explor.* 144, 345–354.
- Greger, M., 2008. Trace elements and radionuclides in edible plants. *Trace Elements as Contaminants and Nutrients*. John Wiley and Sons, Inc., Hoboken, NJ, pp. 121–136.
- Guedes, R., Smaghe, G., Stark, J., Desneux, N., 2016. Pesticide-induced stress in arthropod pests for optimized integrated pest management programs. *Annu. Rev. Entomol.* 61, 43–62.
- Guertin, J., Avakian, C.P., Jacobs, J.A., 2016. *Chromium (VI) Handbook*. CRC press.
- Güngör, E.B.Ö., Bekbölet, M., 2010. Zinc release by humic and fulvic acid as influenced by pH, complexation and DOC sorption. *Geoderma* 159, 131–138.
- Guo, G., Wu, F., Xie, F., Zhang, R., 2012. Spatial distribution and pollution assessment of heavy metals in urban soils from southwest China. *J. Environ. Sci.* 24, 410–418.
- Guoqing, X., Xiuqin, C., Liping, B., Hongtao, Q., Haibo, L., 2019. Absorption, accumulation and distribution of metals and nutrient elements in poplars planted in land amended with composted sewage sludge: a field trial. *Ecotoxicol. Environ. Saf.* 182, 109360.
- Gupta, M., Gupta, S., 2017. An overview of selenium uptake, metabolism, and toxicity in plants. *Front. Plant Sci.* 7, 2074.
- Gupta, S., Nayek, S., Saha, R., Satpati, S., 2008. Assessment of heavy metal accumulation in macrophyte, agricultural soil, and crop plants adjacent to discharge zone of sponge iron factory. *Environ. Geol.* 55, 731–739.
- Gupta, D., Chatterjee, S., Datta, S., Veer, V., Walther, C., 2014. Role of phosphate fertilizers in heavy metal uptake and detoxification of toxic metals. *Chemosphere* 108, 134–144.
- Gupta, D., Chatterjee, S., Datta, S., Voronina, A., Walther, C., 2016. Radionuclides: accumulation and transport in plants. *Reviews of Environmental Contamination and Toxicology*, vol. 241. Springer, pp. 139–160.
- Gupta, D.K., Tiwari, S., Razafindrabe, B., Chatterjee, S., 2017. Arsenic contamination from historical aspects to the present. *Arsenic Contamination in the Environment*. Springer, pp. 1–12.
- Guttikunda, S.K., Jawahar, P., 2014. Atmospheric emissions and pollution from the coal-fired thermal power plants in India. *Atmos. Environ.* 92, 449–460.
- Habibul, N., Hu, Y., Sheng, G.-P., 2016. Microbial fuel cell driving electrokinetic remediation of toxic metal contaminated soils. *J. Hazard. Mater.* 318, 9–14.
- Hajek, A.E., Eilenberg, J., 2018. *Natural Enemies: an Introduction to Biological Control*. Cambridge University Press.
- Halim, M., Majumder, R., Zaman, M., 2015. Paddy soil heavy metal contamination and uptake in rice plants from the adjacent area of Barapukuria coal mine, northwest Bangladesh. *Arab. J. Geosci.* 8, 3391–3401.
- Hashim, M.A., Mukhopadhyay, S., Sahu, J.N., Sengupta, B., 2011. Remediation technologies for heavy metal contaminated groundwater. *J. Environ. Manage.* 92, 2355–2388.
- Hassan, N.U., Mahmood, Q., Waseem, A., Irshad, M., Pervez, A., 2013. Assessment of heavy metals in wheat plants irrigated with contaminated wastewater. *Polish J.*

- Environ. Stud. 22.
- Hawkesford, M.J., De Kok, L.J., 2006. Managing sulphur metabolism in plants. *Plant Cell Environ.* 29, 382–395.
- He, Z.L., Shentu, J., Yang, X.E., 2010. Manganese and Selenium, Trace Elements in Soils. Wiley, London, pp. 481–496.
- Hezbollah, M., Sultana, S., Chakraborty, S., Patwary, M., 2016. Heavy metal contamination of food in a developing country like Bangladesh: an emerging threat to food safety. *J. Toxicol. Environ. Health Sci.* 8, 1–5.
- Hooda, P., 2010. Trace Elements in Soils. John Wiley & Sons.
- Hough, R.L., 2010. Copper and lead. *Trace Elements in Soils.* 441.
- Hseu, Z.-Y., Su, S.-W., Lai, H.-Y., Guo, H.-Y., Chen, T.-C., Chen, Z.-S., 2010. Remediation techniques and heavy metal uptake by different rice varieties in metal-contaminated soils of Taiwan: new aspects for food safety regulation and sustainable agriculture. *Soil Sci. Plant Nutr.* 56, 31–52.
- Hu, L., McBride, M.B., Cheng, H., Wu, J., Shi, J., Xu, J., Wu, L., 2011. Root-induced changes to cadmium speciation in the rhizosphere of two rice (*Oryza sativa* L.) genotypes. *Environ. Res.* 111, 356–361.
- Hu, P., Li, Z., Yuan, C., Ouyang, Y., Zhou, L., Huang, J., Huang, Y., Luo, Y., Christie, P., Wu, L., 2013. Effect of water management on cadmium and arsenic accumulation by rice (*Oryza sativa* L.) with different metal accumulation capacities. *J. Soils Sedim.* 13, 916–924.
- Hu, Y., Cheng, H., Tao, S., 2016. The challenges and solutions for cadmium-contaminated rice in China: a critical review. *Environ. Int.* 92, 515–532.
- Hu, Y., Cheng, H., Tao, S., 2017. Environmental and human health challenges of industrial livestock and poultry farming in China and their mitigation. *Environ. Int.* 107, 111–130.
- Hu, B., Shao, S., Fu, Z., Li, Y., Ni, H., Chen, S., Zhou, Y., Jin, B., Shi, Z., 2019. Identifying heavy metal pollution hot spots in soil-rice systems: a case study in South of Yangtze River Delta, China. *Sci. Total Environ.* 658, 614–625.
- Huang, Z., Pan, X.-D., Wu, P.-G., Han, J.-L., Chen, Q., 2013. Health risk assessment of heavy metals in rice to the population in Zhejiang, China. *PLoS One* 8, e75007.
- Huang, B., Li, Z., Huang, J., Chen, G., Nie, X., Ma, W., Yao, H., Zhen, J., Zeng, G., 2015. Aging effect on the leaching behavior of heavy metals (Cu, Zn, and Cd) in red paddy soil. *Environ. Sci. Pollut. Res. - Int.* 22, 11467–11477.
- Huang, G., Su, X., Rizwan, M.S., Zhu, Y., Hu, H., 2016. Chemical immobilization of Pb, Cu, and Cd by phosphate materials and calcium carbonate in contaminated soils. *Environ. Sci. Pollut. Res. - Int.* 23, 16845–16856.
- Hughes, M.F., Beck, B.D., Chen, Y., Lewis, A.S., Thomas, D.J., 2011. Arsenic exposure and toxicology: a historical perspective. *Toxicol. Sci.* 123, 305–332.
- Huong, N.T.L., Ohtsubo, M., Li, L., Higashi, T., Kanayama, M., Nakano, A., 2008. Heavy metal contamination of soil and rice in wastewater-irrigated paddy field in a suburban area of Hanoi, Vietnam. *Clay science* 13, 205–215.
- Huot, H., Simonnot, M.-O., Morel, J.L., 2015. Pedogenetic trends in soils formed in technogenic parent materials. *Soil Sci.* 180, 182–192.
- Igerase, E., Osifo, P., Ofomaja, A., 2018. Mathematical modelling of Pb²⁺, Cu²⁺, Ni²⁺, Zn²⁺, Cr⁶⁺ and Cd²⁺ ions adsorption from a synthetic acid mine drainage onto chitosan derivative in a packed bed column. *Environ. Technol.* 39, 3203–3220.
- Ihedioha, J.N., Ujam, O.T., Nwuche, C.O., Ekere, N.R., Chime, C.C., 2016. Assessment of heavy metal contamination of rice grains (*Oryza sativa*) and soil from Ada field, Enugu, Nigeria: estimating the human health risk. *Hum. Ecol. Risk Assess. - Int. J.* 22, 1665–1677.
- Ishimaru, Y., Takahashi, R., Bashir, K., Shimo, H., Senoura, T., Sugimoto, K., Ono, K., Yano, M., Ishikawa, S., Arai, T., 2012. Characterizing the role of rice NRAMP5 in manganese, iron and cadmium transport. *Sci. Rep.* 2, 286.
- Islam, M.S., Ahmed, M.K., Habibullah-Al-Mamun, M., 2015. Metal speciation in soil and health risk due to vegetables consumption in Bangladesh. *Environ. Monit. Assess.* 187, 288.
- Islam, M.S., Ahmed, M.K., Habibullah-Al-Mamun, M., 2016. Apportionment of heavy metals in soil and vegetables and associated health risks assessment. *Stoch. Environ. Res. Risk Assess.* 30, 365–377.
- Jadoun, J., Singh, M., 2014. A study of physico-chemical properties & heavy metals in contaminated soils of municipal waste dumpsite at Dholpur. *Int. J. Sci.* 3.
- Jafari, A., Kamarehie, B., Ghaderpoori, M., Khoshnamvand, N., Birjandi, M., 2018. The concentration data of heavy metals in Iranian grown and imported rice and human health hazard assessment. *Data Brief* 16, 453–459.
- Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B.B., Beeregowda, K.N., 2014. Toxicity, mechanism and health effects of some heavy metals. *Interdiscip. Toxicol.* 7, 60–72.
- Jiang, B., Gong, Y., Gao, J., Sun, T., Liu, Y., Oturan, N., Oturan, M.A., 2018. Reductive detoxification of Cr (VI) mediated by environmentally relevant carboxylic acids: state-of-the-art and perspectives. *J. Hazard. Mater.* 365.
- Jiménez, B., Drechsel, P., Koné, D., Bahri, A., 2009. Wastewater, sludge and excreta use in developing countries: an overview. *Wastewater Irrigation and Health.* Routledge, pp. 29–54.
- Jo, I., Koh, M., 2004. Chemical changes in agricultural soils of Korea: data review and suggested countermeasures. *Environ. Geochem. Health* 26, 105–117.
- Jones, G.D., Droz, B., Greve, P., Gottschalk, P., Poffet, D., McGrath, S.P., Seneviratne, S.I., Smith, P., Winkler, L.H., 2017. Selenium deficiency risk predicted to increase under future climate change. *Proc. Natl. Acad. Sci. U. S. A.* 114, 2848–2853.
- Jun-Hui, Z., Hang, M., 2009. Eco-toxicity and metal contamination of paddy soil in an e-wastes recycling area. *J. Hazard. Mater.* 165, 744–750.
- Kendall, B., Dahl, T.W., Anbar, A.D., 2017. The stable isotope geochemistry of molybdenum. *Rev. Mineral. Geochem.* 82, 683–732.
- Khalid, S., Shahid, M., Niazi, N.K., Murtaza, B., Bibi, I., Dumat, C., 2017. A comparison of technologies for remediation of heavy metal contaminated soils. *J. Geochem. Explor.* 182, 247–268.
- Khan, S., Rehman, S., Khan, A.Z., Khan, M.A., Shah, M.T., 2010. Soil and vegetables enrichment with heavy metals from geological sources in Gilgit, northern Pakistan. *Ecotoxicol. Environ. Saf.* 73, 1820–1827.
- Khanam, R., Kumar, A., Nayak, A., Shahid, M., Tripathi, R., Vijayakumar, S., Bhaduri, D., Kumar, U., Mohanty, S., Panneerselvam, P., 2019. Metal (loid)s (As, Hg, Se, Pb and Cd) in paddy soil: bioavailability and potential risk to human health. *Sci. Total Environ.* 699, 134330.
- Khouidi, H., Maatar, Y., Brini, F., Fourati, A., Ammar, N., Masmoudi, K., 2013. Phytoremediation potential of *Arabidopsis thaliana*, expressing ectopically a vacuolar proton pump, for the industrial waste phosphogypsum. *Environ. Sci. Pollut. Res.* 20, 270–280.
- Kibria, M., Islam, M., Alamgir, M., 2012. Influence of waste water irrigation on heavy metal accumulation in soil and plant. *Int. J. Appl. Nat. Sci.* 1, 43–54.
- Kien, C.N., Noi, N.V., Son, L.T., Ngoc, H.M., Tanaka, S., Nishina, T., Iwasaki, K., 2010. Heavy metal contamination of agricultural soils around a chromite mine in Vietnam. *Soil Sci. Plant Nutr.* 56, 344–356.
- Kim, S.A., Guerinot, M.L., 2007. Mining iron: iron uptake and transport in plants. *FEBS Lett.* 581, 2273–2280.
- Komal, T., Mustafa, M., Ali, Z., Kazi, A.G., 2015. Heavy metal uptake and transport in plants. *Heavy Metal Contamination of Soils.* Springer, pp. 181–194.
- Konhauser, K.O., Pecoits, E., Lalonde, S.V., Papineau, D., Nisbet, E.G., Barley, M.E., Arndt, N.T., Zahnle, K., Kamber, B.S., 2009. Oceanic nickel depletion and a methanogen famine before the Great Oxidation Event. *Nature* 458, 750.
- Krishna, A.K., Mohan, K.R., Murthy, N., Periasamy, V., Bipinkumar, G., Manohar, K., Rao, S.S., 2013. Assessment of heavy metal contamination in soils around chromite mining areas, Nuggihalli, Karnataka, India. *Environ. Earth Sci.* 70, 699–708.
- Kumar, P.S., Gunasundari, E., 2018. Bioremediation of heavy metals. *Bioremediation: Applications for Environmental Protection and Management.* Springer, pp. 165–195.
- Kumar, S.S., Kadier, A., Malyan, S.K., Ahmad, A., Bishnoi, N.R., 2017. Phytoremediation and rhizoremediation: uptake, mobilization and sequestration of heavy metals by plants. *Plant-Microbe Interactions in Agro-Ecological Perspectives.* Springer, pp. 367–394.
- Kumarathilaka, P., Seneweera, S., Meharg, A., Bundschuh, J., 2018. Arsenic speciation dynamics in paddy rice soil-water environment: sources, physico-chemical, and biological factors—a review. *Water Res.* 140, 403–414.
- Kumplene, J., Lagerkvist, A., Maurice, C., 2008. Stabilization of As, Cr, Cu, Pb and Zn in soil using amendments—a review. *Waste Manag.* 28, 215–225.
- Kwon, J.C., Nejad, Z.D., Jung, M.C., 2017. Arsenic and heavy metals in paddy soil and polished rice contaminated by mining activities in Korea. *Catena* 148, 92–100.
- Laghlimi, M., Baghdad, B., Hadi, H.E., Bouabdli, A., 2015. Phytoremediation Mechanisms of Heavy Metal Contaminated Soils: a Review.
- Lee, S., An, G., 2009. Over-expression of OsIRT1 leads to increased iron and zinc accumulations in rice. *Plant Cell Environ.* 32, 408–416.
- Lee, S., Jeong, H.J., Kim, S.A., Lee, J., Guerinot, M.L., An, G., 2010. OsZIP5 is a plasma membrane zinc transporter in rice. *Plant Mol. Biol.* 73, 507–517.
- Leng, C.-B., Wang, W., Ye, L., Zhang, X.-C., 2019. Genesis of the late Ordovician Kukaazi Pb-Zn deposit in the western Kunlun orogen, NW China: new insights from in-situ trace elemental compositions of base metal sulfides. *J. Asian Earth Sci.* 184, 103995.
- Leštan, D., Luo, C.-l., Li, X.-d., 2008. The use of chelating agents in the remediation of metal-contaminated soils: a review. *Environ. Pollut.* 153, 3–13.
- Lin, L., Zhou, W., Dai, H., Cao, F., Zhang, G., Wu, F., 2012. Selenium reduces cadmium uptake and mitigates cadmium toxicity in rice. *J. Hazard. Mater.* 235, 343–351.
- Liu, W.-X., Shen, L.-F., Liu, J.-W., Wang, Y.-W., Li, S.-R., 2007. Uptake of toxic heavy metals by rice (*Oryza sativa* L.) cultivated in the agricultural soil near Zhengzhou City, People's Republic of China. *Bull. Environ. Contam. Toxicol.* 79, 209–213.
- Liu, X., Song, Q., Tang, Y., Li, W., Xu, J., Wu, J., Wang, F., Brookes, P.C., 2013a. Human health risk assessment of heavy metals in soil-vegetable system: a multi-medium analysis. *Sci. Total Environ.* 463, 530–540.
- Liu, K.-s., Hao, J.-h., Zeng, Y., Dai, F.-c., Gu, P.-q., 2013b. Neurotoxicity and biomarkers of lead exposure: a review. *Chin. Med. Sci. J.* 28, 178–188.
- Liu, Z., Zhang, Q., Han, T., Ding, Y., Sun, J., Wang, F., Zhu, C., 2016. Heavy metal pollution in a soil-rice system in the Yangtze river region of China. *Int. J. Environ. Res. Public Health* 13, 63.
- Liu, L., Li, W., Song, W., Guo, M., 2018. Remediation techniques for heavy metal-contaminated soils: principles and applicability. *Sci. Total Environ.* 633, 206–219.
- Loganathan, P., Vigneswaran, S., Kandasamy, J., Naidu, R., 2012. Cadmium sorption and desorption in soils: a review. *Crit. Rev. Environ. Sci. Technol.* 42, 489–533.
- Lu, J., Jiang, C., 2001. Antiangiogenic activity of selenium in cancer chemoprevention: metabolite-specific effects. *Nutr. Cancer* 40, 64–73.
- Lu, Y., Song, S., Wang, R., Liu, Z., Meng, J., Sweetman, A.J., Jenkins, A., Ferrier, R.C., Li, H., Luo, W., 2015. Impacts of soil and water pollution on food safety and health risks in China. *Environ. Int.* 77, 5–15.
- Lu, A., Li, B., Li, J., Chen, W., Xu, L., 2018. Heavy metals in paddy soil-rice systems of industrial and township areas from subtropical China: levels, transfer and health risks. *J. Geochem. Explor.* 194, 210–217.
- Luo, L., Ma, Y., Zhang, S., Wei, D., Zhu, Y.-G., 2009. An inventory of trace element inputs to agricultural soils in China. *J. Environ. Manage.* 90, 2524–2530.
- Luo, C., Yang, R., Wang, Y., Li, J., Zhang, G., Li, X., 2012. Influence of agricultural practice on trace metals in soils and vegetation in the water conservation area along the East River (Dongjiang River), South China. *Sci. Total Environ.* 431, 26–32.
- Lwin, C.S., Seo, B.-H., Kim, H.-U., Owens, G., Kim, K.-R., 2018. Application of soil amendments to contaminated soils for heavy metal immobilization and improved soil quality—a critical review. *Soil Sci. Plant Nutr.* 64, 156–167.
- Ma, Y., Hooda, P.S., 2010. Chromium, nickel and cobalt. *Trace Elements in Soils.* pp. 461–480.
- Ma, J.F., Yamaji, N., 2008. Functions and transport of silicon in plants. *Cell. Mol. Life Sci.*

- 65, 3049–3057.
- Ma, Y., Prasad, M., Rajkumar, M., Freitas, H., 2011. Plant growth promoting rhizobacteria and endophytes accelerate phytoremediation of metalliferous soils. *Biotechnol. Adv.* 29, 248–258.
- Ma, Y., Oliveira, R.S., Nai, F., Rajkumar, M., Luo, Y., Rocha, I., Freitas, H., 2015. The hyperaccumulator *Sedum plumbizincicola* harbors metal-resistant endophytic bacteria that improve its phytoextraction capacity in multi-metal contaminated soil. *J. Environ. Manage.* 156, 62–69.
- Mahar, A., Wang, P., Ali, A., Awasthi, M.K., Lahori, A.H., Wang, Q., Li, R., Zhang, Z., 2016. Challenges and opportunities in the phytoremediation of heavy metals contaminated soils: a review. *Ecotoxicol. Environ. Saf.* 126, 111–121.
- Majed, N., Real, M., Isreq, H., Akter, M., Azam, H.M., 2016. Food adulteration and biomagnification of environmental contaminants: a comprehensive risk framework for Bangladesh. *Front. Environ. Sci.* 4, 34.
- Makino, T., 2007. Heavy Metal Pollution of Soil and a New Approach to Its Remediation: Research Experiences in Japan. Food and Fertilizer Technology Center.
- Makino, T., Luo, Y., Wu, L., Sakurai, Y., Maejima, Y., Akahane, I., Arai, T., 2010. Heavy metal pollution of soil and risk alleviation methods based on soil chemistry (< Special Issue > International symposium: Challenges to Soil Degradation Towards Sustaining Life and Environment, Tokyo Metropolitan University Symposium Series No. 2, 2009). *Pedologist* 53, 38–49.
- Mani, A., Sankaranarayanan, K., 2018. Heavy metal and mineral element-induced abiotic stress in rice plant. *Rice Crop: Current Developments.* 149.
- Marques, A.P., Rangel, A.O., Castro, P.M., 2009. Remediation of heavy metal contaminated soils: phytoremediation as a potentially promising clean-up technology. *Crit. Rev. Environ. Sci. Technol.* 39, 622–654.
- Marques, A.P., Rangel, A.O., Castro, P.M., 2011. Remediation of heavy metal contaminated soils: an overview of site remediation techniques. *Crit. Rev. Environ. Sci. Technol.* 41, 879–914.
- Mateo-Sagasta, J., Raschid-Sally, L., Thebo, A., 2015. Global wastewater and sludge production, treatment and use. *Wastewater*. Springer, pp. 15–38.
- Matović, V., Buha, A., Đukić-Čosić, D., Bulat, Z., 2015. Insight into the oxidative stress induced by lead and/or cadmium in blood, liver and kidneys. *Food Chem. Toxicol.* 78, 130–140.
- Mazej, D., Osvald, J., Stibilj, V., 2008. Selenium species in leaves of chicory, dandelion, lamb's lettuce and parsley. *Food Chem.* 107, 75–83.
- McCann, C.M., 2012. The Use of Natural Mn Oxide-Containing Wastes as a Contaminated Land Remediation Strategy and their Effects on Soil Microbial Functioning. Newcastle University.
- Meharg, A.A., Rahman, M.M., 2003. Arsenic contamination of Bangladesh paddy field soils: implications for rice contribution to arsenic consumption. *Environ. Sci. Technol.* 37, 229–234.
- Menezes-Filho, J.A., Paes, C.R., Pontes, Â.Md.C., Moreira, J.C., Sarcinelli, P.N., Mergler, D., 2009. High levels of hair manganese in children living in the vicinity of a ferro-manganese alloy production plant. *Neurotoxicology* 30, 1207–1213.
- Milner, M.J., Seamon, J., Craft, E., Kochian, L.V., 2013. Transport properties of members of the ZIP family in plants and their role in Zn and Mn homeostasis. *J. Exp. Bot.* 64, 369–381.
- Minnikova, T., Denisova, T., Mandzhieva, S., Kolesnikov, S., Minkina, T., Chaplygin, V., Burachevskaya, M., Sushkova, S., Bauer, T., 2015. Assessing the effect of heavy metals from the Novochebarkassk power station emissions on the biological activity of soils in the adjacent areas. *J. Geochem. Explor.* 174, 70–78.
- Mishra, A., Tripathi, B.D., 2008. Heavy metal contamination of soil, and bioaccumulation in vegetables irrigated with treated waste water in the tropical city of Varanasi, India. *Toxicol. Environ. Chem.* 90, 861–871.
- Moreno-Jiménez, E., Esteban, E., Peñalosa, J.M., 2012. The fate of arsenic in soil-plant systems. *Reviews of Environmental Contamination and Toxicology*. Springer, pp. 1–37.
- Murtaza, G., Ghafoor, A., Qadir, M., Owens, G., Aziz, M., Zia, M., 2010. Disposal and use of sewage on agricultural lands in Pakistan: a review. *Pedosphere* 20, 23–34.
- Musa, J., Mustapha, H., Bala, J., Ibrahim, Y., Akos, M., Daniel, E., Oguche, F., Kutu, I., 2017a. Heavy metals in agricultural soils in Nigeria: a review, arid zone journal of engineering. *Technol. Environ.* 13, 593–603.
- Musa, J., Mustapha, H., Bala, J., Ibrahim, Y., Akos, M., Daniel, E., Oguche, F., Kutu, I., 2017b. Heavy metals in agricultural soils in Nigeria: a review, arid zone journal of engineering. *Technol. Environ.* 13, 593.
- Mushtaq, F., Mat, R., Ani, F.N., 2014. A review on microwave assisted pyrolysis of coal and biomass for fuel production. *Renew. Sustain. Energy Rev.* 39, 555–574.
- Nair, A.T., Ahammed, M.M., 2015. The reuse of water treatment sludge as a coagulant for post-treatment of UASB reactor treating urban wastewater. *J. Clean. Prod.* 96, 272–281.
- Navarro-Alarcón, M., Cabrera-Vique, C., 2008. Selenium in food and the human body: a review. *Sci. Total Environ.* 400, 115–141.
- Nayak, A., Raja, R., Rao, K., Shukla, A., Mohanty, S., Shahid, M., Tripathi, R., Panda, B., Bhattacharyya, P., Kumar, A., 2015. Effect of fly ash application on soil microbial response and heavy metal accumulation in soil and rice plant. *Ecotoxicol. Environ. Saf.* 114, 257–262.
- Nejad, Z.D., Jung, M.C., Kim, K.-H., 2018. Remediation of soils contaminated with heavy metals with an emphasis on immobilization technology. *Environ. Geochem. Health* 40, 927–953.
- Nieder, R., Benbi, D.K., Reichl, F.X., 2018. Role of potentially toxic elements in soils. *Soil Components and Human Health*. Springer, pp. 375–450.
- Nogawa, K., Kobayashi, E., Okubo, Y., Suwazono, Y., 2004. Environmental cadmium exposure, adverse effects and preventive measures in Japan. *Biomaterials* 17, 581–587.
- Nordstrom, D.K., 2011. Hydrogeochemical processes governing the origin, transport and fate of major and trace elements from mine wastes and mineralized rock to surface waters. *Appl. Geochem.* 26, 1777–1791.
- Nzengue, Y., Candéas, S.M., Sauvaigo, S., Douki, T., Favier, A., Rachidi, W., Guiraud, P., 2011. The toxicity redox mechanisms of cadmium alone or together with copper and zinc homeostasis alteration: its redox biomarkers. *J. Trace Elem. Med. Biol.* 25, 171–180.
- Nziguheba, G., Smolders, E., 2008. Inputs of trace elements in agricultural soils via phosphate fertilizers in European countries. *Sci. Total Environ.* 390, 53–57.
- Olowokere, B., Oyibo, O., Oyerinde, G., 2018. Heavy metals concentration in dumpsites at Gwagwalada, Abuja: implications on sustainable environmental management. *Greener J. Soil Sci. Plant Nutr.* 5, 001–004.
- Padmavathamma, P.K., Li, L.Y., 2007. Phytoremediation technology: hyper-accumulation metals in plants. *Water Air Soil Pollut.* 184, 105–126.
- Pahade, V., Sharma, A., 2015. Manganese removal by low cost adsorbent from synthetic wastewater—a review. *Int. J. Eng. Res.* 4, 111–114.
- Palaniappan, S., Annadurai, K., 2018. Organic Farming Theory & Practice. Scientific publishers.
- Palansooriya, K.N., Shaheen, S.M., Chen, S.S., Tsang, D.C., Hashimoto, Y., Hou, D., Bolan, N.S., Rinklebe, J., Ok, Y.S., 2020. Soil amendments for immobilization of potentially toxic elements in contaminated soils: a critical review. *Environ. Int.* 134, 105046.
- Pan, Y., 2015. Speciation of Trace Metals and Their Uptake by Rice in Paddy Soils. Wageningen University.
- Parth, V., Murthy, N., Saxena, P.R., 2011. Assessment of heavy metal contamination in soil around hazardous waste disposal sites in Hyderabad city (India): natural and anthropogenic implications. *J. Environ. Res. Manage.* 2, 027–034.
- Pathak, C., Chopra, A., Kumar, V., Srivastava, S., 2010. Heavy metals contamination in waste-water irrigated agricultural soil near Bindal river, Dehradun, India. *Pollut. Res.* 29, 583â.
- Patrick, L., Toxicity, Lead, 2006. A review of the literature. Part I: exposure, evaluation, and treatment. *Altern. Med. Rev.* 11.
- Pettygrove, G.S., 2018. Irrigation with Reclaimed Municipal wastewater—A Guidance Manual. CRC Press.
- Pilon-Smits, E.A., Quinn, C.F., 2010. Selenium metabolism in plants. *Cell Biology of Metals and Nutrients*. Springer, pp. 225–241.
- Pinto, E., Aguiar, A.A., Ferreira, I.M., 2014. Influence of soil chemistry and plant physiology in the phytoremediation of Cu, Mn, and Zn. *Crit. Rev. Plant Sci.* 33, 351–373.
- Pinto, A., De Varennes, A., Fonseca, R., Teixeira, D.M., 2015. Phytoremediation of soils contaminated with heavy metals: techniques and strategies. *Phytoremediation*. Springer, pp. 133–155.
- Pinto, A., de Varennes, A., Dias, C., Lopes, M., 2018. Microbial-assisted phytoremediation: a convenient use of plant and microbes to clean up soils. *Phytoremediation*. Springer, pp. 21–87.
- Potysz, A., Grybos, M., Kierczak, J., Guibaud, G., Fondaneche, P., Lens, P.N., van Hullebusch, E.D., 2017. Metal mobilization from metallurgical wastes by soil organic acids. *Chemosphere* 178, 197–211.
- Prasad, R., Shivay, Y.S., Kumar, D., 2016. Interactions of zinc with other nutrients in soils and plants—a review. *Indian J. Fertil.* 12, 16–26.
- Prasanna, R., 2017. Bioprospecting of Neem and Hibiscus against Corrosion Caused by Soil Microbes on Zinc and Its Alloys Buried in Soil.
- Qadir, M., Wichelns, D., Raschid-Sally, L., McCormick, P.G., Drechsel, P., Bahri, A., Minhas, P., 2010. The challenges of wastewater irrigation in developing countries. *Agric. Water Manage.* 97, 561–568.
- Qin, C., Luo, C., Chen, Y., Shen, Z., 2012. Spatial-based assessment of metal contamination in agricultural soils near an abandoned copper mine of Eastern China. *Bull. Environ. Contam. Toxicol.* 89, 113–118.
- Rahman, M.A., Hasegawa, H., Rahman, M.M., Rahman, M.A., Miah, M., 2007. Accumulation of arsenic in tissues of rice plant (*Oryza sativa* L.) and its distribution in fractions of rice grain. *Chemosphere* 69, 942–948.
- Rahman, M.A., Hasegawa, H., Lim, R.P., 2012. Bioaccumulation, biotransformation and trophic transfer of arsenic in the aquatic food chain. *Environ. Res.* 116, 118–135.
- Rahman, M.A., Rahman, M.M., Reichman, S.M., Lim, R.P., Naidu, R., 2014. Heavy metals in Australian grown and imported rice and vegetables on sale in Australia: health hazard. *Ecotoxicol. Environ. Saf.* 100, 53–60.
- Rajasulochana, P., Preethy, V., 2016. Comparison on efficiency of various techniques in treatment of waste and sewage water—A comprehensive review. *Resour. Technol.* 2, 175–184.
- Ravenscroft, P., Brammer, H., Richards, K., 2009. Arsenic Pollution: a Global Synthesis. John Wiley & Sons.
- Rengel, Z., 2000. Manganese uptake and transport. *Metal Ions in Biological Systems: Manganese and its Role in Biological Processes*, vol. 37 57.
- Renkema, H., Koopmans, A., Kersbergen, L., Kikkert, J., Hale, B., Berkelaar, E., 2012. The effect of transpiration on selenium uptake and mobility in durum wheat and spring canola. *Plant Soil* 354, 239–250.
- Rinklebe, J., Shaheen, S.M., 2017. Redox chemistry of nickel in soils and sediments: a review. *Chemosphere* 179, 265–278.
- Rinklebe, J., Antoniadis, V., Shaheen, S.M., Rosche, O., Altermann, M., 2019. Health risk assessment of potentially toxic elements in soils along the Central Elbe River, Germany. *Environ. Int.* 126, 76–88.
- Rodríguez, O., Padilla, I., Tayibi, H., López-Delgado, A., 2012. Concerns on liquid mercury and mercury-containing wastes: a review of the treatment technologies for the safe storage. *J. Environ. Manage.* 101, 197–205.
- Rogan, N., Serafimovski, T., Dolenc, M., Tasev, G., Dolenc, T., 2009. Heavy metal contamination of paddy soils and rice (*Oryza sativa* L.) from Kočani Field (Macedonia). *Environ. Geochem. Health* 31, 439–451.
- Roy, M., Giri, A.K., Dutta, S., Mukherjee, P., 2015. Integrated phytobal remediation for sustainable management of arsenic in soil and water. *Environ. Int.* 75, 180–198.
- Roychowdhury, T., Tokunaga, H., Ando, M., 2003. Survey of arsenic and other heavy

- metals in food composites and drinking water and estimation of dietary intake by the villagers from an arsenic-affected area of West Bengal, India. *Sci. Total Environ.* 308, 15–35.
- Rudrabasher, T.S.M., 2016. The Prevalence of Heavy Metals (Arsenic) in Soils around Kuala Lumpur City, Malaysia 6, 104–107.
- Saha, R., Nandi, R., Saha, B., 2011. Sources and toxicity of hexavalent chromium. *J. Coord. Chem.* 64, 1782–1806.
- Sahu, S., Ajmal, P., Bhangare, R., Tiwari, M., Pandit, G., 2014. Natural radioactivity assessment of a phosphate fertilizer plant area. *J. Radiat. Res. Appl. Sci.* 7, 123–128.
- Salmnikow, K., Zhitkovich, A., 2007. Genetic and epigenetic mechanisms in metal carcinogenesis and cocarcinogenesis: nickel, arsenic, and chromium. *Chem. Res. Toxicol.* 21, 28–44.
- Sarbas, B., 2013. Mn Manganese: Natural Occurrence. Minerals (Native Metal, Solid Solution, Silicide, and Carbide. Sulfides and Related Compounds. Halogenides and Oxyhalogenides. Oxides of Type MO). Springer Science & Business Media.
- Sarwar, N., Imran, M., Shaheen, M.R., Ishaque, W., Kamran, M.A., Matloob, A., Rehman, Hussain, S., 2017. Phytoremediation strategies for soils contaminated with heavy metals: modifications and future perspectives. *Chemosphere* 171, 710–721.
- Sasaki, A., Yamaji, N., Yokosho, K., Ma, J.F., 2012. Nramp5 is a major transporter responsible for manganese and cadmium uptake in rice. *Plant Cell* 24, 2155–2167.
- Satpathy, D., Reddy, M.V., Dhal, S.P., 2014. Risk assessment of heavy metals contamination in paddy soil, plants, and grains (*Oryza sativa* L.) at the East Coast of India. *Biomed Res. Int.* 2014.
- Sattari, S., Van Ittersum, M., Giller, K., Zhang, F., Bouwman, A., 2014. Key role of China and its agriculture in global sustainable phosphorus management. *Environ. Res. Lett.* 9, 054003.
- Savci, S., 2012. An agricultural pollutant: chemical fertilizer. *Int. J. Environ. Sci. Dev.* 3, 73.
- Scerri, E., 2019. The Periodic Table: Its Story and Its Significance. Oxford University Press.
- Seck, P.A., Diagne, A., Mohanty, S., Wopereis, M.C., 2012. Crops that feed the world 7: rice. *Food Secur.* 4, 7–24.
- Sessitsch, A., Kuffner, M., Kidd, P., Vangronsveld, J., Wenzel, W.W., Fallmann, K., Puschenreiter, M., 2013. The role of plant-associated bacteria in the mobilization and phytoextraction of trace elements in contaminated soils. *Soil Biol. Biochem.* 60, 182–194.
- Shaheen, S.M., Tsadilas, C.D., Rinklebe, J., 2013. A review of the distribution coefficients of trace elements in soils: influence of sorption system, element characteristics, and soil colloidal properties. *Adv. Colloid Interface Sci.* 201, 43–56.
- Shahid, M., Xiong, T., Masood, N., Leveque, T., Quenea, K., Austruy, A., Foucault, Y., Dumat, C., 2014. Influence of plant species and phosphorus amendments on metal speciation and bioavailability in a smelter impacted soil: a case study of food-chain contamination. *J. Soils Sedim.* 14, 655–665.
- Shahid, M., Shamshad, S., Rafiq, M., Khalid, S., Bibi, I., Niazi, N.K., Dumat, C., Rashid, M.I., 2017a. Chromium speciation, bioavailability, uptake, toxicity and detoxification in soil-plant system: a review. *Chemosphere* 178, 513–533.
- Shahid, M., Dumat, C., Khalid, S., Schreck, E., Xiong, T., Niazi, N.K., 2017b. Foliar heavy metal uptake, toxicity and detoxification in plants: a comparison of foliar and root metal uptake. *J. Hazard. Mater.* 325, 36–58.
- Shahid, M., Niazi, N.K., Khalid, S., Murtaza, B., Bibi, I., Rashid, M.I., 2018. A critical review of selenium biogeochemical behavior in soil-plant system with an inference to human health. *Environ. Pollut.* 234, 915–934.
- Shi, J., Yu, X., Zhang, M., Lu, S., Wu, W., Wu, J., Xu, J., 2011. Potential risks of copper, zinc, and cadmium pollution due to pig manure application in a soil-rice system under intensive farming: a case study of Nanhu, China. *J. Environ. Qual.* 40, 1695–1704.
- Shraim, A.M., 2017. Rice is a potential dietary source of not only arsenic but also other toxic elements like lead and chromium. *Arab. J. Chem.* 10, S3434–S3443.
- Simmons, R.W., Noble, A., Pongsakul, P., Sukreeyapongse, O., Chinabut, N., 2008. Analysis of field-moist Cd contaminated paddy soils during rice grain fill allows reliable prediction of grain Cd levels. *Plant Soil* 302, 125–137.
- Singh, R., Agrawal, M., 2010. Variations in heavy metal accumulation, growth and yield of rice plants grown at different sewage sludge amendment rates. *Ecotoxicol. Environ. Saf.* 73, 632–641.
- Singh, R., Gautam, N., Mishra, A., Gupta, R., 2011a. Heavy metals and living systems: an overview. *Indian J. Pharmacol.* 43, 246.
- Singh, J., Upadhyay, S.K., Pathak, R.K., Gupta, V., 2011b. Accumulation of heavy metals in soil and paddy crop (*Oryza sativa*), irrigated with water of Ramgarh Lake, Gorakhpur, UP, India. *Toxicol. Environ. Chem.* 93, 462–473.
- Singh, M., Garg, V., Gautam, Y., Kumar, A., 2014. Soil to Grain Transfer Factors of Heavy Metals in Rice and Health Risk Analysis in the Vicinity of Narora Atomic Power Station (NAPS), Narora, India.
- Singh, R., Singh, S., Parihar, P., Singh, V.P., Prasad, S.M., 2015. Arsenic contamination, consequences and remediation techniques: a review. *Ecotoxicol. Environ. Saf.* 112, 247–270.
- Singh, H., Verma, A., Kumar, M., Sharma, R., Gupta, R., Kaur, M., Negi, M., Sharma, S., 2017. Phytoremediation: a green technology to clean up the sites with low and moderate level of heavy metals. *Austin Biochem.* 2, 1012.
- Sivakumar, D., Kandaswamy, A., Gomathi, V., Rajeshwaran, R., Murugan, N., 2014. Bioremediation studies on reduction of heavy metals toxicity. *Pollut. Res.* 33, 553–558.
- Smičiklas, I., Smiljanić, S., Perić-Grujić, A., Šljivić-Ivanović, M., Mitrić, M., Antonović, D., 2014. Effect of acid treatment on red mud properties with implications on Ni (II) sorption and stability. *Chem. Eng. J.* 242, 27–35.
- Soares, M.A., Quina, M.J., Quinta-Ferreira, R.M., 2015. Immobilisation of lead and zinc in contaminated soil using compost derived from industrial eggshell. *J. Environ. Manage.* 164, 137–145.
- Soetan, K., Olaiya, C., Oyewole, O., 2010. The importance of mineral elements for humans, domestic animals and plants—a review. *Afr. J. Food Sci.* 4, 200–222.
- Soliman, M., Potlakayala, S., Millar, D., Weeden, H., Bogush, D., Deguchi, M., Rudrabhatla, S., 2019. Comparing a review of heavy metal uptake and their toxicity on plant and human health. *Int. J. Plant Anim. Environ. Sci.* 9, 182–189.
- Song, W.-Y., Park, J., Eisenach, C., Maeshima, M., Lee, Y., Martinoia, E., 2014. ABC transporters and heavy metals. *Plant ABC Transporters*. Springer, pp. 1–17.
- Stroud, J.L., Norton, G.J., Islam, M.R., Dasgupta, T., White, R.P., Price, A.H., Meharg, A.A., McGrath, S.P., Zhao, F.-J., 2011. The dynamics of arsenic in four paddy fields in the Bengal delta. *Environ. Pollut.* 159, 947–953.
- Sun, W., Huang, B., Zhao, Y., Shi, X., Darilek, J.L., Deng, X., Wang, H., Zou, Z., 2009. Spatial variability of soil selenium as affected by geologic and pedogenic processes and its effect on ecosystem and human health. *Geochim. J.* 43, 217–225.
- Sun, G.-X., Liu, X., Williams, P.N., Zhu, Y.-G., 2010. Distribution and translocation of selenium from soil to grain and its speciation in paddy rice (*Oryza sativa* L.). *Environ. Sci. Technol.* 44, 6706–6711.
- Sun, H.-J., Rathinasabapathi, B., Wu, B., Luo, J., Pu, L.-P., Ma, L.Q., 2014. Arsenic and selenium toxicity and their interactive effects in humans. *Environ. Int.* 69, 148–158.
- Tack, F.M., Meers, E., 2010. Assisted phytoextraction: helping plants to help us. *Elements* 6, 383–388.
- Tak, H.I., Ahmad, F., Babalola, O.O., 2013. Advances in the application of plant growth-promoting rhizobacteria in phytoremediation of heavy metals. *Reviews of Environmental Contamination and Toxicology*, vol. 223. Springer, pp. 33–52.
- Tan, H., Zhang, G., Heaney, P.J., Webb, S.M., Burgos, W.D., 2010. Characterization of manganese oxide precipitates from Appalachian coal mine drainage treatment systems. *Appl. Geochem.* 25, 389–399.
- Tangahu, B.V., Abdullah, S., Rozaimah, S., Basri, H., Idris, M., Anuar, N., Mukhlisin, M., 2011. A review on heavy metals (As, Pb, and Hg) uptake by plants through phytoremediation. *Int. J. Chem. Eng.* 2011.
- Tao, X.-Q., Shen, D.-S., Shentu, J.-L., Long, Y.-Y., Feng, Y.-J., Shen, C.-C., 2015. Bioaccessibility and health risk of heavy metals in ash from the incineration of different e-waste residues. *Environ. Sci. Pollut. Res. - Int.* 22, 3558–3569.
- Tariq, S., 2001. Environmental Geochemistry of Surface and Sub Surface Water and Soil in Peshawar Basin of NWFP Pakistan. Unpublished PhD thesis. University of Peshawar.
- Tariq, S.R., Rashid, N., 2012. Multivariate analysis of metal levels in paddy soil, rice plants, and rice grains: a case study from Shakargarh, Pakistan. *J. Chem.* 2013.
- Teklay, A., 2016. Physiological effect of chromium exposure: a review. *Int. J. Food Sci. Nutr. Diet.* 7, 1–11.
- Thakur, S., Singh, L., Ab Wahid, Z., Siddiqui, M.F., Atnaw, S.M., Din, M.F.M., 2016. Plant-driven removal of heavy metals from soil: uptake, translocation, tolerance mechanism, challenges, and future perspectives. *Environ. Monit. Assess.* 188, 206.
- Timmer, C.P., 2014. Food security in a sea and the Pacific: the rapidly changing role of rice. *Asia Pac. Policy Stud.* 1, 73–90.
- Tinggi, U., 2003. Essentiality and toxicity of selenium and its status in Australia: a review. *Toxicol. Lett.* 137, 103–110.
- Tiwari, M., Sharma, D., Dwivedi, S., Singh, M., Tripathi, R.D., Trivedi, P.K., 2014. Expression in a *rabidopsis* and cellular localization reveal involvement of rice NRAMP, OsNRAMP 1, in arsenic transport and tolerance. *Plant Cell Environ.* 37, 140–152.
- Tripathi, A., Misra, D., 2012. A study of physico-chemical properties and heavy metals in contaminated soils of municipal waste dumpsites at Allahabad, India. *Int. J. Environ. Sci.* 2, 2024.
- Tsukada, H., Hasegawa, H., Takeda, A., Hisamatsu, S., 2007. Concentrations of major and trace elements in polished rice and paddy soils collected in Aomori, Japan. *J. Radioanal. Nucl. Chem.* 273, 199–203.
- Tue, N.M., Goto, A., Takahashi, S., Itai, T., Asante, K.A., Kunisue, T., Tanabe, S., 2016. Release of chlorinated, brominated and mixed halogenated dioxin-related compounds to soils from open burning of e-waste in Agbogbloshie (Accra, Ghana). *J. Hazard. Mater.* 302, 151–157.
- Uddin, I., Bano, A., Masood, S., 2015. Chromium toxicity tolerance of *Solanum nigrum* L. and *Parthenium hysterophorus* L. plants with reference to ion pattern, antioxidant activity and root exudation. *Ecotoxicol. Environ. Saf.* 113, 271–278.
- Uraguchi, S., Fujiwara, T., 2012. Cadmium transport and tolerance in rice: perspectives for reducing grain cadmium accumulation. *Rice* 5, 5.
- Uraguchi, S., Fujiwara, T., 2013. Rice breaks ground for cadmium-free cereals. *Curr. Opin. Plant Biol.* 16, 328–334.
- Verkleij, J.A., Golan-Goldhirsh, A., Antosiewicz, D.M., Schwitzguébel, J.-P., Schröder, P., 2009. Dualities in plant tolerance to pollutants and their uptake and translocation to the upper plant parts. *Environ. Exp. Bot.* 67, 10–22.
- Vinceti, M., Maraldi, T., Bergomi, M., Malagoli, C., 2009. Risk of chronic low-dose selenium overexposure in humans: insights from epidemiology and biochemistry. *Rev. Environ. Health* 24, 231–248.
- Violante, A., 2013. Elucidating mechanisms of competitive sorption at the mineral/water interface. *Advances in Agronomy*. Elsevier, pp. 111–176.
- Violante, A., Cozzolino, V., Perelomov, L., Caporale, A., Pigna, M., 2010. Mobility and bioavailability of heavy metals and metalloids in soil environments. *J. Soil Sci. Plant Nutr.* 10, 268–292.
- Vithanage, M., Rajapaksha, A.U., Oze, C., Rajakaruna, N., Dissanayake, C., 2014. Metal release from serpentine soils in Sri Lanka. *Environ. Monit. Assess.* 186, 3415–3429.
- Vocciante, M., Caretta, A., Bua, L., Bagatin, R., Ferro, S., 2016. Enhancements in ElectroKinetic Remediation Technology: environmental assessment in comparison with other configurations and consolidated solutions. *Chem. Eng. J.* 289, 123–134.
- Vodyanitskii, Y.N., 2009. Mineralogy and geochemistry of manganese: a review of publications. *Eurasian Soil Sci.* 42, 1170–1178.

- Wan, X., Lei, M., Chen, T., 2016. Cost-benefit calculation of phytoremediation technology for heavy-metal-contaminated soil. *Sci. Total Environ.* 563, 796–802.
- Wang, B., Xie, Z., Chen, J., Jiang, J., Su, Q., 2008. Effects of field application of phosphate fertilizers on the availability and uptake of lead, zinc and cadmium by cabbage (*Brassica chinensis* L.) in a mining tailing contaminated soil. *J. Environ. Sci.* 20, 1109–1117.
- Wang, P., Sun, G., Jia, Y., Meharg, A.A., Zhu, Y., 2014. A review on completing arsenic biogeochemical cycle: microbial volatilization of arsines in environment. *J. Environ. Sci.* 26, 371–381.
- Wang, X., Zeng, X., Chuanping, L., Li, F., Xu, X., Lv, Y., 2016. Heavy metal contaminations in soil-rice system: source identification in relation to a sulfur-rich coal burning power plant in Northern Guangdong Province, China. *Environ. Monit. Assess.* 188, 460.
- Wang, Z., Li, J., Li, Y., 2017. Using reclaimed water for agricultural and landscape irrigation in China: a review. *Irrig. Drain.* 66, 672–686.
- Wang, W., Mauleon, R., Hu, Z., Chebotarov, D., Tai, S., Wu, Z., Li, M., Zheng, T., Fuentes, R.R., Zhang, F., 2018. Genomic variation in 3,010 diverse accessions of Asian cultivated rice. *Nature* 557, 43.
- Weston, L.A., Ryan, P.R., Watt, M., 2012. Mechanisms for cellular transport and release of allelochemicals from plant roots into the rhizosphere. *J. Exp. Bot.* 63, 3445–3454.
- White, R.E., 2013. Principles and Practice of Soil Science: the Soil as a Natural Resource. John Wiley & Sons.
- Wilson, S.C., Lockwood, P.V., Ashley, P.M., Tighe, M., 2010. The chemistry and behaviour of antimony in the soil environment with comparisons to arsenic: a critical review. *Environ. Pollut.* 158, 1169–1181.
- Winkel, L.H., Johnson, C.A., Lenz, M., Grundl, T., Leupin, O.X., Amini, M., Charlet, L., 2011. Environmental Selenium Research: From Microscopic Processes to Global Understanding. ACS Publications.
- Winkel, L., Vriens, B., Jones, G., Schneider, L., Pilon-Smits, E., Bañuelos, G., 2015. Selenium cycling across soil-plant-atmosphere interfaces: a critical review. *Nutrients* 7, 4199–4239.
- Wortman, S.E., Lovell, S.T., 2013. Environmental challenges threatening the growth of urban agriculture in the United States. *J. Environ. Qual.* 42, 1283–1294.
- Wu, F., Fu, Z., Liu, B., Mo, C., Chen, B., Corns, W., Liao, H., 2011. Health risk associated with dietary co-exposure to high levels of antimony and arsenic in the world's largest antimony mine area. *Sci. Total Environ.* 409, 3344–3351.
- Wuana, R.A., Okieimen, F.E., 2011. Heavy metals in contaminated soils: a review of sources, chemistry, risks and best available strategies for remediation. *ISRN Ecol.* 2011.
- Xia, S., Song, Z., Jeyakumar, P., Shaheen, S.M., Rinklebe, J., Ok, Y.S., Bolan, N., Wang, H., 2019. A critical review on bioremediation technologies for Cr (VI)-contaminated soils and wastewater. *Crit. Rev. Environ. Sci. Technol.* 49, 1027–1078.
- Yadav, I.C., Devi, N.L., Syed, J.H., Cheng, Z., Li, J., Zhang, G., Jones, K.C., 2015. Current status of persistent organic pesticides residues in air, water, and soil, and their possible effect on neighboring countries: a comprehensive review of India. *Sci. Total Environ.* 511, 123–137.
- Yadav, P., Singh, B., Garg, V., Mor, S., Pulhani, V., 2017. Bioaccumulation and health risks of heavy metals associated with consumption of rice grains from croplands in Northern India. *Hum. Ecol. Risk Assess.: Int. J.* 23, 14–27.
- Yadav, K.K., Gupta, N., Kumar, A., Reece, L.M., Singh, N., Rezanian, S., Khan, S.A., 2018. Mechanistic understanding and holistic approach of phytoremediation: a review on application and future prospects. *Ecol. Eng.* 120, 274–298.
- Yang, X., Feng, Y., He, Z., Stoffella, P.J., 2005. Molecular mechanisms of heavy metal hyperaccumulation and phytoremediation. *J. Trace Elem. Med. Biol.* 18, 339–353.
- Yang, S., Sheng, G., Tan, X., Hu, J., Du, J., Montavon, G., Wang, X., 2011. Determination of Ni (II) uptake mechanisms on morденite surfaces: a combined macroscopic and microscopic approach. *Geochim. Cosmochim. Acta* 75, 6520–6534.
- Yang, G., Zhang, G., Wang, H., 2015. Current state of sludge production, management, treatment and disposal in China. *Water Res.* 78, 60–73.
- Yang, Z., Zhang, Z., Chai, L., Wang, Y., Liu, Y., Xiao, R., 2016. Bioremediation of heavy metal-contaminated soils using *Burkholderia* sp. Z-90. *Journal of Hazardous Materials* 301, 145–152.
- Yao, Z., Li, J., Xie, H., Yu, C., 2012. Review on remediation technologies of soil contaminated by heavy metals. *Procedia Environ. Sci.* 16, 722–729.
- Yap, D., Adezrian, J., Khairiah, J., Ismail, B., Ahmad-Mahir, R., 2009. The uptake of heavy metals by paddy plants (*Oryza sativa*) in Kota Marudu, Sabah, Malaysia. *Am. J. Agric. Environ. Sci.* 6, 16–19.
- Yin, K., Ahamed, A., Lisak, G., 2018. Environmental perspectives of recycling various combustion ashes in cement production—a review. *Waste Manag.* 78, 401–416.
- Zain, M.E., 2011. Impact of mycotoxins on humans and animals. *J. Saudi Chem. Soc.* 15, 129–144.
- Zalewska, M., 2012. Response of perennial ryegrass (*Lolium perenne* L.) to soil contamination with zinc. *J. Elem.* 17.
- Zeng, H., 2009. Selenium as an essential micronutrient: roles in cell cycle and apoptosis. *Molecules* 14, 1263–1278.
- Zeng, H., Uthus, E.O., Combs, G.F., 2005. Mechanistic aspects of the interaction between selenium and arsenic. *J. Inorg. Biochem.* 99, 1269–1274.
- Zeng, F., Ali, S., Zhang, H., Ouyang, Y., Qiu, B., Wu, F., Zhang, G., 2011. The influence of pH and organic matter content in paddy soil on heavy metal availability and their uptake by rice plants. *Environ. Pollut.* 159, 84–91.
- Zeng, F., Wei, W., Li, M., Huang, R., Yang, F., Duan, Y., 2015. Heavy metal contamination in rice-producing soils of Hunan province, China and potential health risks. *Int. J. Environ. Res. Public Health* 12, 15584–15593.
- Zhang, J., 2011. Fate and Transport of Arsenic in Construction and Demolition Debris Landfills and Soils Underneath. University of Florida.
- Zhang, Y., Zhang, H.-W., Zhen-Cheng, S., Zhang, C.-G., 2008. Soil microbial characteristics under long-term heavy metal stress: a case study in Zhangshi wastewater irrigation area, Shenyang. *Pedosphere* 18, 1–10.
- Zhang, W., Jiang, F., Ou, J., 2011. Global pesticide consumption and pollution: with China as a focus. *Proc. Int. Acad. Ecol. Environ. Sci.* 1, 125.
- Zhang, H., Feng, X., Zhu, J., Sapkota, A., Meng, B., Yao, H., Qin, H., Larssen, T., 2012a. Selenium in soil inhibits mercury uptake and translocation in rice (*Oryza sativa* L.). *Environ. Sci. Technol.* 46, 10040–10046.
- Zhang, W.-H., Ying-Xin, W., Simonnot, M., 2012b. Soil contamination due to e-waste disposal and recycling activities: a review with special focus on China. *Pedosphere* 22, 434–455.
- Zhang, C., Wu, P., Tang, C., Han, Z., Sun, J., 2013. Assessment of arsenic distribution in paddy soil and rice plants of a typical karst basin affected by acid mine drainage in Southwest China. *Environ. Pollut.* 2, 27.
- Zhang, H., Feng, X., Jiang, C., Li, Q., Liu, Y., Gu, C., Shang, L., Li, P., Lin, Y., Larssen, T., 2014. Understanding the paradox of selenium contamination in mercury mining areas: high soil content and low accumulation in rice. *Environ. Pollut.* 188, 27–36.
- Zhang, S.-Y., Zhao, F.-J., Su, G.-X., Su, J.-Q., Yang, X.-R., Li, H., Zhu, Y.-G., 2015. Diversity and abundance of arsenic biotransformation genes in paddy soils from southern China. *Environ. Sci. Technol.* 49, 4138–4146.
- Zhang, W., Sheng, R., Zhang, M., Xiong, G., Hou, H., Li, S., Wei, W., 2018. Effects of continuous manure application on methanogenic and methanotrophic communities and methane production potentials in rice paddy soil. *Agric. Ecosyst. Environ.* 258, 121–128.
- Zhang, Y.-y., Lv, J.-w., Dong, X.-j., Fang, Q., Tan, W.-f., Wu, X.-y., Deng, Q.-w., 2019. Influence on Uranium (VI) migration in soil by iron and manganese salts of humic acid: mechanism and behavior. *Environ. Pollut.* 256, 113369.
- Zhao, F.J., Ma, J.F., Meharg, A., McGrath, S., 2009a. Arsenic uptake and metabolism in plants. *New Phytol.* 181, 777–794.
- Zhao, K., Zhang, W., Zhou, L., Liu, X., Xu, J., Huang, P., 2009b. Modeling transfer of heavy metals in soil-rice system and their risk assessment in paddy fields. *Environ. Earth Sci.* 59, 519–527.
- Zhao, K., Liu, X., Xu, J., Selim, H., 2010. Heavy metal contaminations in a soil-rice system: identification of spatial dependence in relation to soil properties of paddy fields. *J. Hazard. Mater.* 181, 778–787.
- Zhao, F.-J., Ma, Y., Zhu, Y.-G., Tang, Z., McGrath, S.P., 2014a. Soil contamination in China: current status and mitigation strategies. *Environ. Sci. Technol.* 49, 750–759.
- Zhao, Y., Yan, Z., Qin, J., Xiao, Z., 2014b. Effects of long-term cattle manure application on soil properties and soil heavy metals in corn seed production in Northwest China. *Environ. Sci. Pollut. Res. - Int.* 21, 7586–7595.
- Zhao, F.-J., Ma, Y., Zhu, Y.-G., Tang, Z., McGrath, S.P., 2015a. Soil contamination in China: current status and mitigation strategies. *Environ. Sci. Technol.* 49, 750–759.
- Zhao, K., Fu, W., Ye, Z., Zhang, C., 2015b. Contamination and spatial variation of heavy metals in the soil-rice system in Nanxun County, Southeastern China. *Int. J. Environ. Res. Public Health* 12, 1577–1594.
- Zhengfu, B., Inyang, H.I., Daniels, J.L., Frank, O., Struthers, S., 2010. Environmental issues from coal mining and their solutions. *Min. Sci. Technol.* 20, 215–223.
- Ziarati, P., Shirkan, F., Mostafidi, M., Zahedi, M., 2018. An overview of the heavy metal contamination in milk and dairy products. *Acta Sci. Pharm. Sci.* 2, 8–21.
- Zwolak, I., 2019. The role of selenium in arsenic and cadmium toxicity: an updated review of scientific literature. *Biol. Trace Elem. Res.* 1–20.
- Zwolak, I., Zaporowska, H., 2012. Selenium interactions and toxicity: a review. *Cell Biol. Toxicol.* 28, 31–46.